

Sr. No.

Total No. of Printed pages : 23

10498

(DO NOT OPEN THIS QUESTION BOOKLET BEFORE TIME OR UNTIL YOU ARE ASKED TO DO SO)

PG-EE-2013
Five Year Mathematics (Hons.)

Code

B

Time : 1¼ hours

Max. Marks : 100

Total Questions : 100

Roll No. _____ (in figure) _____ (in words)

Name _____ Father's Name _____

Mother's Name _____ Date of Examination : _____

(Signature of the candidate)

(Signature of the Invigilator)

CANDIDATES MUST READ THE FOLLOWING INFORMATION / INSTRUCTIONS BEFORE STARTING THE QUESTION PAPER.

1. All questions are compulsory and carry equal marks.
2. The candidate must return the Question book-let as well as OMR answer-sheet to the Invigilator concerned before leaving the Examination Hall, failing which a case of use of unfair-means / misbehaviour will be registered against him / her, in addition to lodging of an FIR with the police. Further the answer-sheet of such a candidate will not be evaluated.
3. In case there is any discrepancy in any question(s) in the Question Booklet, the same may be brought to the notice of the Controller of Examinations in writing **within two hours** after the test is over. No such complaint(s) will be entertained thereafter.
4. The candidate **MUST NOT** do any rough work or writing in the OMR Answer-Sheet. Rough work, if any, may be done in the question book-let itself. Answers **MUST NOT** be ticked in the Question book-let.
5. Use only blue or black **BALL POINT PEN** of good quality in the OMR Answer-Sheet.
6. There will be no negative marking. Each correct answer will be awarded one full mark. Cutting, erasing, overwriting and more than one answer in OMR Answer-Sheet will be treated as incorrect answer.
7. **BEFORE ANSWERING THE QUESTIONS, THE CANDIDATES SHOULD ENSURE THAT THEY HAVE BEEN SUPPLIED CORRECT AND COMPLETE BOOK-LET. COMPLAINTS, IF ANY, REGARDING MISPRINTING ETC. WILL NOT BE ENTERTAINED 30 MINUTES AFTER STARTING OF THE EXAMINATION.**



45/104
22/7/13

Question No.	Questions
1.	Negation of $p \rightarrow q$ is (1) $\sim p \vee q$ (2) $p \wedge (\sim q)$ (3) $\sim q \rightarrow \sim p$ (4) $p \vee (\sim q)$
2.	Five observations are given as 25, 25, 25, 25 and 25. The mean and standard deviation of these observations are respectively (1) 5 and 5 (2) 25 and 5 (3) 25 and 25 (4) 25 and 0
3.	If the median of 11 observations is 20 and if the observations greater than the median are increased by 5, then the median of the new data will be (1) 20 (2) 25 (3) $25 + \frac{20}{11}$ (4) $25 - \frac{20}{11}$
4.	An event is called a simple event if it has (1) only two sample points of a sample space (2) more than two sample points of a sample space (3) only one sample point of a sample space (4) No sample point of a sample space
5.	If A and B are two mutually exclusive events, then which of the following may not be true (1) occurrence of any one of them excludes the occurrence of the other event. (2) A and B cannot occur simultaneously (3) A and B are disjoint (4) A and B are equally likely

Question No.	Questions
6.	Which of the following probabilities are not consistently defined ? (1) $P(A) = 0.5, P(B) = 0.7, P(A \cup B) = 0.6$ (2) $P(A) = 0.5, P(B) = 0.7, P(A \cap B) = 0.4$ (3) $P(A) = 0.5, P(B) = 0.4, P(A \cup B) = 0.8$ (4) $P(A) = 0.6, P(B) = 0.7, P(A \cup B) = 0.8$
7.	The probability that a student will pass the final examination in both English and Hindi is 0.5 and the probability of passing neither is 0.2. If the probability of passing the English examination is 0.75, the probability of passing the Hindi examination is (1) 1 (2) 0.55 (3) 0.05 (4) 0.45
8.	The number of all possible matrices of order 3×3 with each entry 1 or 2 is (1) 18 (2) 27 (3) 256 (4) 512
9.	Which of the following is not true for a square matrix A ? (1) A can be expressed as the sum of a symmetric and a skew symmetric matrix (2) If A is skew symmetric matrix, then all its diagonal elements are zero (3) $A + A'$ is a skew symmetric matrix (4) A is symmetric if $A' = A$.

Question No.	Questions
10.	If $A = \begin{bmatrix} \cos 2\alpha & -\sin 2\alpha \\ \sin 2\alpha & \cos 2\alpha \end{bmatrix}$, then $A + A' = I$, if the value of α is
	(1) $\frac{\pi}{6}$ (2) $\frac{\pi}{3}$ (3) π (4) $\frac{3\pi}{2}$
11.	IQ of a person is given by the formula $IQ = \frac{MA}{CA} \times 100$, where MA is mental age and CA is chronological age. If $84 \leq IQ \leq 144$ for a group of 12 years old children, the range of their mental age is
	(1) $7 \leq MA \leq 12$ (2) $10.08 \leq MA \leq 17.28$ (3) $0 \leq MA \leq 12$ (4) $0 \leq MA \leq 7$
12.	Number of different signals that can be generated by arranging at least 3 flags in order (one below the other) on a vertical staff, if five different flags are available, is
	(1) 15 (2) 125 (3) 243 (4) 300
13.	The least positive integer n for which ${}^{n-1}C_3 + {}^{n-1}C_4 < {}^nC_5$ is
	(1) 4 (2) 5 (3) 9 (4) 10

Question No.	Questions
14.	<p>If letters of the word RADHIK are arranged in all positive ways and are written out as in a dictionary, then the word RADHIK appears at serial number</p> <p>(1) 600 (2) 601 (3) 120 (4) 121</p>
15.	<p>For a positive integer n, the value of ${}^n C_0 - {}^n C_1 + {}^n C_2 - \dots + (-1)^n \cdot {}^n C_n$ is</p> <p>(1) 0 (2) 1 (3) -1 (4) 2^n</p>
16.	<p>The remainder when 2^{300} is divided by 9 is</p> <p>(1) 0 (2) 1 (3) 2 (4) 8</p>
17.	<p>If the length of sides of a right triangle are in A. P., then the sines of acute angles of the triangle are</p> <p>(1) $\frac{1}{3}, \frac{2}{3}$ (2) $\sqrt{\frac{3}{5}}, \sqrt{\frac{2}{3}}$ (3) $\sqrt{\frac{1}{3}}, \sqrt{\frac{2}{3}}$ (4) $\frac{3}{5}, \frac{4}{5}$</p>
18.	<p>If the sum of the series $3 + \frac{3}{x} + \frac{9}{x^2} + \frac{27}{x^3} + \dots$ is finite, then</p> <p>(1) $-3 < x < 3$ (2) $-1 < x < 1$ (3) $x > 9$ (4) $x > 3$</p>

Question No.	Questions
19.	<p>If three points $(h, 0)$, (a, b) and $(0, k)$ lie on a line, then</p> <p>(1) $\frac{a}{h} - \frac{b}{k} = 1$ (2) $\frac{a}{h} + \frac{b}{k} = 1$</p> <p>(3) $\frac{b}{k} - \frac{a}{h} = 1$ (4) $\frac{a}{h} + \frac{b}{k} = -1$</p>
20.	<p>The value (s) of k for which the line $(k - 3)x - (4 - k^2)y + k^2 - 7k + 6 = 0$ is parallel to y-axis is</p> <p>(1) 3 (2) ± 3 (3) 6, 1 (4) ± 2</p>
21.	<p>If A, B, C are three non-empty sets such that $A \cap B = \phi$, $B \cap C = \phi$, then</p> <p>(1) $A = C$ (2) $A \subset C$</p> <p>(3) $C \subset A$ (4) None of these</p>
22.	<p>Two finite sets have m and n elements respectively. The total number of subsets of second set is 112 more than the total number of subsets of the first set. The values of m and n respectively are</p> <p>(1) 7, 8 (2) 4, 7 (3) 6, 8 (4) 3, 7</p>
23.	<p>The set of all second elements of the ordered pairs in a relation R from a set A to set B is called the</p> <p>(1) domain of the relation R</p> <p>(2) Range of the relation R</p> <p>(3) co-domain of the relation R</p> <p>(4) None of these</p>

Question No.	Questions
24.	<p>Let $R = \{(x, y) : x, y \in A, x + y = 7\}$, where $A = \{1, 2, 3, 4, 5, 6, 7\}$, then</p> <p>(1) R is symmetric but not reflexive and not transitive</p> <p>(2) R is an equivalence relation</p> <p>(3) R is reflexive, symmetric but not transitive</p> <p>(4) R is not reflexive, not symmetric but is transitive</p>
25.	<p>Domain and range respectively of the function $f(x) = \sqrt{4 - x^2}$ are</p> <p>(1) $\{x : -2 \leq x \leq 2\}, \{x : -2 \leq x \leq 2\}$</p> <p>(2) $\{x : -2 \leq x \leq 2\}, \{x : 0 \leq x \leq 2\}$</p> <p>(3) $\{x : 0 \leq x \leq 2\}, \{x : -2 \leq x \leq 2\}$</p> <p>(4) $\{x : 0 \leq x \leq 2\}, \{x : 0 \leq x \leq 2\}$</p>
26.	<p>Let $A = \{1, 2, 3, 4\}$, $B = \{1, 5, 9, 11, 15, 16\}$ and $f = \{(1, 5), (2, 9), (3, 1), (4, 5), (2, 11)\}$.</p> <p>Which of the following is true ?</p> <p>(1) f is a relation from A to B</p> <p>(2) f is a function from A to B</p> <p>(3) f is a relation from B to A</p> <p>(4) f is a function from B to A</p>
27.	<p>The function $f : \mathbb{N} \rightarrow \mathbb{N}$ given by $f(x) = 3x$ is</p> <p>(1) one-one and onto (2) one-one but not onto</p> <p>(3) onto but not one-one (4) Neither one-one nor onto</p>

Question No.	Questions
28.	<p>Consider a binary operation $*$ on N defined as $a * b = a^2 + b^2$. Choose the correct answer</p> <p>(1) $*$ is both associative and commutative (2) $*$ is associative but not commutative (3) $*$ is commutative but not associative (4) $*$ is neither commutative nor associative</p>
29.	<p>If $\cos 32^\circ = m$ and $\cos x = 2m^2 - 1$; α, β are the values of x between 0° and 360°, then</p> <p>(1) $\alpha + \beta = 180^\circ$ (2) $\beta - \alpha = 200^\circ$ (3) $\beta = 4\alpha + 40^\circ$ (4) $\beta = 5\alpha - 20^\circ$</p>
30.	<p>Which of the following is true for</p> $\tan(x+y) = \frac{\tan x + \tan y}{1 - \tan x \tan y} ?$ <p>(1) Angles x, y are odd multiple of $\frac{\pi}{2}$ and $(x+y)$ is multiple of π (2) Angles x, y are multiple of π and $(x+y)$ is odd multiple of $\frac{\pi}{2}$ (3) None of the angles x, y and $x+y$ is an odd multiple of $\frac{\pi}{2}$ (4) None of the angles x, y and $x+y$ is a multiple of π</p>
31.	<p>If $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$, the equation of the plane through $(3, 4, -1)$ which is parallel to the plane $2x - 3y + 5z + 7 = 0$ is</p> <p>(1) $\vec{r} \cdot (2\hat{i} - 3\hat{j} + 5\hat{k}) + 11 = 0$ (2) $\vec{r} \cdot (3\hat{i} + 4\hat{j} - \hat{k}) + 11 = 0$ (3) $\vec{r} \cdot (3\hat{i} - 4\hat{j} - \hat{k}) + 7 = 0$ (4) $\vec{r} \cdot (2\hat{i} - 3\hat{j} + 5\hat{k}) - 7 = 0$</p>

Question No.	Questions
32.	The constants in a linear programming problem are (1) linear (2) quadratic (3) cubic (4) biquadratic
33.	The common region determined by all the constants including non-negative constraints of a linear programming problem is called the (1) optimal solution (2) feasible solution (3) infeasible solution (4) unbounded solution
34.	The corner points of the feasible region determined by the following system of linear inequalities : $2x + y \leq 10, x + 3y \leq 15; x, y \geq 0$ are $(0, 0), (5, 0), (3, 4)$ and $(0, 5)$. Let $Z = px + qy$, where $p, q > 0$. Condition on p and q so that the maximum of Z occurs at both $(3, 4)$ and $(0, 5)$ is (1) $p = q$ (2) $p = 2q$ (3) $q = 3p$ (4) $p = 3q$
35.	If A and B be two events such that $P(A) = 0.4, P(A \cup B) = 0.8$. If A and B are independent events, then the probability P(B) is (1) $\frac{2}{5}$ (2) $\frac{3}{5}$ (3) $\frac{1}{5}$ (4) $\frac{2}{3}$

Question No.	Questions
36.	If A and B are two events such that $0 < P(B) < 1$, then (1) $P(A \bar{B}) + P(\bar{A} \bar{B}) = 1$ (2) $P(A B) + P(A \bar{B}) = 1$ (3) $P(\bar{A} B) + P(A \bar{B}) = 1$ (4) None of these
37.	If the standard deviation of the binomial distribution $(q + p)^{16}$ is 2, then mean of the distribution is (1) 6 (2) 8 (3) 10 (4) 12
38.	A fair coin is tossed repeatedly. If head and tail appear alternatively on first 5 tosses, then the probability that head appears on the sixth toss is (1) $\frac{1}{2}$ (2) $\frac{1}{32}$ (3) $\frac{1}{64}$ (4) $\frac{5}{64}$
39.	A and B toss a coin alternatively till one of them gets a head and wins the game. If A begins the game, the probability that B wins the game is (1) $\frac{1}{2}$ (2) $\frac{1}{3}$ (3) $\frac{1}{4}$ (4) $\frac{2}{3}$
40.	Posteriori probability for an event is obtained using (1) Additive law of probability (2) Multiplication theorem of probability (3) Bayes' theorem (4) Classical definition of probability

Question No.	Questions
41.	<p>Let $h(x) = \min \{x, x^2\}$ for every real number x. Then</p> <p>(1) h is continuous for all x</p> <p>(2) h is differentiable for all x</p> <p>(3) $h'(x) = 0$ for all $x > 1$</p> <p>(4) h is differentiable at two values of x, that is, 0 and 1</p>
42.	<p>Let a function f be defined by $f(x) = \frac{x - x }{x}$ for $x \neq 0$ and $f(0) = 2$. Then f is</p> <p>(1) continuous nowhere</p> <p>(2) continuous everywhere</p> <p>(3) continuous for all x except at $x = 1$</p> <p>(4) continuous for all x except at $x = 0$</p>
43.	<p>$\frac{d}{dx} [\tan^{-1}(\sec x + \tan x)]$ is equal to</p> <p>(1) 0</p> <p>(2) $\sec x - \tan x$</p> <p>(3) $\frac{1}{2}$</p> <p>(4) 2</p>
44.	<p>If $x = \log t$ and $y = t^2 - 1$, then $\frac{d^2y}{dx^2}$ at $t = 2$ is</p> <p>(1) 8</p> <p>(2) 16</p> <p>(3) 4</p> <p>(4) 2</p>

Question No.	Questions
45.	<p>If $y = \sin^{-1} \left(\frac{1-x^2}{1+x^2} \right)$, $0 < x < 1$; then $\frac{dy}{dx}$ is equal to</p> <p>(1) $\frac{2}{\sqrt{1-x^2}}$ (2) $\frac{-2}{\sqrt{1-x^2}}$</p> <p>(3) $\frac{2}{1+x^2}$ (4) $\frac{-2}{1+x^2}$</p>
46.	<p>Let A and B be two points on the graph of function $y = f(x)$ corresponding to $x = a$ and $x = b$. If Lagrange's mean value theorem is applicable over the interval $[a, b]$, then there exists at least one point on the graph between A and B, the tangent at which is parallel to</p> <p>(1) x-axis (2) y-axis</p> <p>(3) the chord AB (4) line $y = x$</p>
47.	<p>The rate of change of the volume of a sphere with respect to its radius r at $r = 6$ cm is</p> <p>(1) 144π (2) 48π</p> <p>(3) 432π (4) 12π</p>
48.	<p>The points on the curve $y = x^3$ at which the slope of the tangent is equal to the y-coordinate of the point are</p> <p>(1) $(0, 0), (1, 3)$ (2) $(0, 0), (2, 8)$</p> <p>(3) $(0, 0), (3, 27)$ (4) $(0, 0), (4, 48)$</p>

Question No.	Questions
49.	<p>The point on the curve $x^2 = 2y$ in the second quadrant which is nearest to the point $(0, 5)$ is</p> <p>(1) $(-2, 2)$ (2) $(-2\sqrt{2}, 4)$ (3) $(-1, \frac{1}{2})$ (4) $(-\sqrt{2}, 1)$</p>
50.	<p>If $\frac{d}{dx} f(x) = \sin 2x - 4e^{3x}$ such that $f(0) = \frac{7}{6}$, then $f(x)$ is</p> <p>(1) $-\frac{1}{2} \cos 2x - \frac{4}{3} e^{3x} + 3$ (2) $\cos 2x - 4e^{3x} - \frac{11}{6}$ (3) $\frac{1}{2} \cos 2x - \frac{4e^{3x}}{3} - 3$ (4) $-\frac{1}{2} \cos 2x - \frac{4}{3} e^{3x} - 3$</p>
51.	<p>Let the generator of a double-napped right circular cone be inclined to its vertical axis at an angle α. A plane cuts the nappe (other than the vertex) of the cone making an angle β with the vertical axis of the cone. The section so obtained on this intersection is parabola if</p> <p>(1) $\beta = 90^\circ$ (2) $\alpha < \beta < 90^\circ$ (3) $\beta = \alpha$ (4) $0 \leq \beta < \alpha$</p>
52.	<p>In an ellipse, the distance between the foci is 6 and minor axis is 8, then the eccentricity is</p> <p>(1) $\frac{3}{4}$ (2) $\frac{3}{5}$ (3) $\frac{4}{5}$ (4) $\frac{2}{3}$</p>

Question No.	Questions
53.	Length of latus rectum of the hyperbola $\frac{y^2}{9} - \frac{x^2}{27} = 1$ is (1) 18 (2) $2\sqrt{3}$ (3) 6 (4) $\frac{2}{3}$
54.	Ratio in which the line segment joining the points (4, 8, 10) and (6, 10, -8) is divided by the xz-plane is (1) 2 : 3 externally (2) 2 : 3 internally (3) 4 : 5 externally (4) 5 : 4 internally
55.	If the origin is the centroid of a triangle PQR and the co-ordinates of its two vertices P and Q are (-4, 2, 6) and (-4, -16, -10) respectively, then the co-ordinates of the vertex R are (1) $\left(-\frac{8}{3}, -\frac{14}{3}, -\frac{4}{3}\right)$ (2) (-8, -14, -4) (3) $\left(\frac{8}{3}, \frac{14}{3}, \frac{4}{3}\right)$ (4) (8, 14, 4)
56.	$\lim_{x \rightarrow 0} \frac{\sqrt{1 - \cos 2x}}{\sqrt{2x}}$ (1) exists and it equals to 1 (2) exists and it equals to -1 (3) exists and it equals to 0 (4) does not exist

Question No.	Questions
57.	<p>If $\lim_{x \rightarrow 0} \frac{\sin px}{\tan 3x} = 4$, then the value of p is</p> <p>(1) $\frac{3}{4}$ (2) $\frac{4}{3}$ (3) 12 (4) 4</p>
58.	<p>The derivative of an even function is always</p> <p>(1) an odd function (2) an even function</p> <p>(3) does not exist (4) None of these</p>
59.	<p>If $f'(3) = 2$, then $\lim_{h \rightarrow 0} \frac{f(3+h^2) - f(3-h^2)}{2h^2}$ is</p> <p>(1) 1 (2) 2 (3) 0 (4) $\frac{1}{2}$</p>
60.	<p>Which of the following sentences is not a statement ?</p> <p>(1) There are 35 days in a month</p> <p>(2) The sum of 5 and 7 is greater than 10</p> <p>(3) Mathematics is difficult</p> <p>(4) All real numbers are complex numbers</p>
61.	<p>Choose the correct answer :</p> <p>$\int \frac{20x^{19} + 20^x \log_e 20}{x^{20} + 20^x} dx$ equals</p> <p>(1) $x^{20} + 20^x + c$ (2) $\log \left(\frac{1}{x^{20} + 20^x} \right) + c$</p> <p>(3) $\log (20x^{19} + 20^x \log_e 20) + c$ (4) $\log (x^{20} + 20^x) + c$</p>

Question No.	Questions
62.	<p>The value of $\sqrt{2} \int \frac{\sin x}{\sin\left(x - \frac{\pi}{4}\right)} dx$ is</p> <p>(1) $x + \log \left \cos\left(x - \frac{\pi}{4}\right) \right + c$</p> <p>(2) $x - \log \left \sin\left(x - \frac{\pi}{4}\right) \right + c$</p> <p>(3) $x + \log \left \sin\left(x - \frac{\pi}{4}\right) \right + c$</p> <p>(4) $x - \log \left \cos\left(x - \frac{\pi}{4}\right) \right + c$</p>
63.	<p>The function $f(x) = \int \frac{x-2}{x^2-7x+12} dx$</p> <p>(1) decreases on \mathbb{R}</p> <p>(2) increases on $\mathbb{R} - (2, 3)$</p> <p>(3) increases on $(2, 3) \cup (4, \infty)$</p> <p>(4) $(2, \infty)$</p>
64.	<p>$f(x) = \int \frac{dx}{\sin^4 x}$ is a</p> <p>(1) polynomial of degree 3 in $\cot x$</p> <p>(2) polynomial of degree 4 in $\cot x$</p> <p>(3) polynomial of degree 4 in $\operatorname{cosec} x$</p> <p>(4) polynomial of degree 3 in $\operatorname{cosec} x$</p>

Question No.	Questions
70.	Let $f(x) = \int_1^x e^{-t^2/2} (1-t^2) dt$, then f has (1) maximum at $x=0$ (2) maximum at $x=-1$ (3) maximum at $x=-1$ (4) no critical point
71.	The degree of the differential equation $\left(\frac{d^2y}{dx^2}\right)^{3/2} - \left(\frac{dy}{dx}\right)^{1/2} - 4 = 0$ is (1) 6 (2) 4 (3) 3 (4) 2
72.	The number of arbitrary constants in the particular solution of a differential equation of second order is (1) 3 (2) 2 (3) 1 (4) 0
73.	The general solution of the differential equation $\frac{dy}{dx} = e^{x-y}$ is (1) $e^x - e^y = c$ (2) $e^x - e^{-y} = c$ (3) $e^{-x} - e^y = c$ (4) $e^x + e^y = c$
74.	Direction cosines of the vector $\hat{i} + \hat{j} - 2\hat{k}$ are (1) $(1, 1, -2)$ (2) $\left(\frac{1}{2}, \frac{1}{2}, -1\right)$ (3) $\left(\frac{1}{2\sqrt{2}}, \frac{1}{2\sqrt{2}}, -\frac{1}{\sqrt{2}}\right)$ (4) $\left(\frac{1}{\sqrt{6}}, \frac{1}{\sqrt{6}}, -\frac{2}{\sqrt{6}}\right)$

Question No.	Questions
75.	Projection of vector $2\hat{i} + 3\hat{j} + 2\hat{k}$ on the vector $\hat{i} + 2\hat{j} + \hat{k}$ is (1) $\frac{2\sqrt{15}}{3}$ (2) $\frac{5}{3}\sqrt{6}$ (3) 10 (4) 6
76.	If \vec{a} and \vec{b} are two unit vectors and θ is the angle between them. Then $\vec{a} - \vec{b}$ is a unit vector if (1) $\theta = \frac{\pi}{4}$ (2) $\theta = \frac{\pi}{3}$ (3) $\theta = \frac{\pi}{2}$ (4) $\theta = \frac{2\pi}{3}$
77.	$(\vec{a} + \vec{b}) \cdot (\vec{a} + \vec{b}) = \vec{a} ^2 + \vec{b} ^2$ if and only if (1) $\vec{a} = \vec{b}$ (2) \vec{a} is parallel to \vec{b} (3) \vec{a}, \vec{b} are perpendicular (4) $\vec{a} + \vec{b} = 0$
78.	If a line makes angles $90^\circ, 135^\circ, 45^\circ$ with the x, y and z-axis respectively, then its direction cosines are (1) $0, \frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}$ (2) $0, \frac{1}{2}, \frac{\sqrt{3}}{2}$ (3) 1, 0, 0 (4) $0, -\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}$
79.	Distance of the point (0, 0, 0) from the plane $3x - 4y + 12z = 3$ is (1) 0 (2) $\frac{1}{3}$ (3) $\frac{3}{13}$ (4) $\frac{3}{11}$

Question No.	Questions
80.	<p>The angle between the lines $2x = 3y = -z$ and $6x = -y = -4z$ is</p> <p>(1) $\frac{\pi}{4}$ (2) $\frac{\pi}{6}$ (3) 0 (4) $\frac{\pi}{2}$</p>
81.	<p>For any real numbers x and y, $\cos x = \cos y$ implies</p> <p>(1) $x = n\pi + (-1)^n y$, where $n \in \mathbb{Z}$</p> <p>(2) $x = n\pi \pm y$, where $n \in \mathbb{Z}$</p> <p>(3) $x = n\pi + y$, where $n \in \mathbb{Z}$</p> <p>(4) $x = (2n + 1)\frac{\pi}{2} + y$, where $n \in \mathbb{Z}$</p>
82.	<p>If the roots of the quadratic equation $x^2 + px + q = 0$ are $\tan 30^\circ$ and $\tan 15^\circ$, then the value of $2 + q - p$ is</p> <p>(1) 0 (2) 1 (3) 2 (4) 3</p>
83.	<p>If $\cos^{-1} x + \cos^{-1} y = \frac{2\pi}{3}$, then $\sin^{-1} x + \sin^{-1} y$ is equal to</p> <p>(1) $\frac{2\pi}{3}$ (2) $\frac{\pi}{3}$</p> <p>(3) $\frac{\pi}{6}$ (4) π</p>
84.	<p>Principal value of $\cot^{-1} \left(-\frac{1}{\sqrt{3}} \right)$ is</p> <p>(1) $\frac{2\pi}{3}$ (2) $\frac{\pi}{3}$</p> <p>(3) $-\frac{2\pi}{3}$ (4) $-\frac{\pi}{3}$</p>

Question No.	Questions
85.	$\tan^{-1} \left(\frac{x}{y} \right) - \tan^{-1} \frac{x-y}{x+y}$ is equal to (1) $\frac{\pi}{2}$ (2) $\frac{\pi}{3}$ (3) $\frac{\pi}{4}$ (4) $-\frac{3\pi}{4}$
86.	$3 \cos^{-1} x - \pi x - \frac{\pi}{2} = 0$ has (1) one solution (2) one and only one solution (3) no solution (4) more than one solution
87.	A set S is said to be an inductive set if (1) $x+1 \in S$ implies $x \in S$ and $1 \notin S$ (2) $x+1 \in S$ implies $x \in S$ and $1 \in S$ (3) $x \in S$ implies $1 \in S$ (4) $1 \in S$ and $x+1 \in S$ whenever $x \in S$
88.	If $\left(\frac{1+i}{1-i} \right)^x = 1$ and n is any positive integer then (1) $x = 2n$ (2) $x = 4n + 1$ (3) $x = 2n + 1$ (4) $x = 4n$
89.	The argument of complex number $\frac{1}{1+i}$ is (1) $\frac{\pi}{4}$ (2) $-\frac{\pi}{4}$ (3) $\frac{\pi}{2}$ (4) $-\frac{\pi}{2}$

Question No.	Questions
96.	<p>If A is an invertible matrix of order 3 and $\det(A) = 3$, then $\det(A^{-1})$ is equal to</p> <p>(1) $\frac{1}{3}$ (2) 3 (3) 9 (4) 0</p>
97.	<p>The value of k for which the system of equations</p> $x + ky - 3z = 0$ $3x + ky - 2z = 0$ $2x + 3y - 4z = 0$ <p>has a non-trivial solution is</p> <p>(1) $\frac{21}{10}$ (2) 2 (3) $\frac{31}{10}$ (4) 4</p>
98.	<p>Minor of an element of a determinant of order 4 is a determinant of order</p> <p>(1) 4 (2) 3 (3) 2 (4) 1</p>
99.	<p>Let A and B are square matrices of the same order with $A = 3$ and $B = -5$, then AB is</p> <p>(1) $\frac{5}{3}$ (2) 15 (3) -15 (4) None of these</p>
100.	<p>Matrix equation of a system of linear equations is $AX = B$ and A is a singular matrix, then the system of equations is called inconsistent if</p> <p>(1) $(\text{adj } A)B = 0$ (2) $\text{Adj } A = 0$</p> <p>(3) $B = 0$ (4) $(\text{adj } A)B \neq 0$</p>