

Found sealed and opened for verification during OMR evaluation at 1:20 PM

Total No. of Printed Pages : 13

(DO NOT OPEN THIS QUESTION BOOKLET BEFORE TIME OR UNTIL YOU ARE ASKED TO DO SO)

PG-EE-2016

SUBJECT : Mathematics Hons. Five Year

A

Sr. No. 10101

Time : 1½ Hours

Max. Marks : 100

Total Questions : 100

Roll No. (in figures) _____ (in words) _____

Name _____ Father's Name _____

Mother's Name _____ Date of Examination _____

(Signature of the Candidate)

(Signature of the Invigilator)

CANDIDATES MUST READ THE FOLLOWING INFORMATION/INSTRUCTIONS BEFORE STARTING THE QUESTION PAPER.

1. All questions are compulsory.
2. The candidates **must return** the question booklet as well as OMR Answer-Sheet to the Invigilator concerned before leaving the Examination Hall, failing which a case of use of unfair-means/misbehaviour will be registered against him/her, in addition to lodging of an FIR with the police. Further the answer-sheet of such a candidate will not be evaluated.
3. In case there is any discrepancy in any question(s) in the Question Booklet, the same may be brought to the notice of the Controller of Examinations in writing **within two hours** after the test is over. No such complaint(s) will be entertained thereafter.
4. The candidate **must not** do any rough work or writing in the OMR Answer-Sheet. Rough work, if any, may be done in the question booklet itself. Answers **must not** be ticked in the question booklet.
5. **There will be no negative marking. Each correct answer will be awarded one full mark. Cutting, erasing, overwriting and more than one answer in OMR Answer-Sheet will be treated as incorrect answer.**
6. Use only **Black** or **Blue Ball Point Pen** of good quality in the OMR Answer-Sheet.
7. *Before answering the questions, the candidates should ensure that they have been supplied correct and complete booklet. Complaints, if any, regarding misprinting etc. will not be entertained 30 minutes after starting of the examination.*

PG-EE-2016/(Maths Hons Five Year)/(A)

1. If two sets A and B are having 99 elements in common, then the number of elements common to each of the sets $A \times B$ and $B \times A$ are :
- (1) 99^2 (2) 18 (3) 2^{99} (4) 100
2. If $f(x) = 1 - \frac{1}{x}$, then the value of $f\left[f\left(\frac{1}{x}\right)\right]$ is :
- (1) $\frac{1}{x} - 1$ (2) $\frac{x}{1-x}$ (3) $\frac{x-1}{x}$ (4) $\frac{x}{x-1}$
3. The function $f(x) = \log\left(x + \sqrt{x^2 + 1}\right)$ is :
- (1) an even function (2) an odd function
(3) a periodic function (4) none of these
4. The value of $\sin A \sin(60^\circ + A) \sin(60^\circ - A)$ is equal to :
- (1) $\sin 3A$ (2) $\sin \frac{3A}{2}$ (3) $\sin \frac{3A}{4}$ (4) $\sin \frac{4A}{3}$
5. If $\cos \theta = \frac{1}{2}\left(x + \frac{1}{x}\right)$, then $\frac{1}{2}\left(x^2 + \frac{1}{x^2}\right)$ is equal to :
- (1) $\cos 2\theta$ (2) $\sin 2\theta$ (3) $\sec 2\theta$ (4) $\tan 2\theta$
6. If $y = \sin^2 \theta + \operatorname{cosec}^2 \theta$, $\theta \neq 0$, then :
- (1) $y > 2$ (2) $y \leq 2$ (3) $y \geq -2$ (4) $y = 0$
7. The value of $\frac{\cos 12^\circ - \sin 12^\circ}{\cos 12^\circ + \sin 12^\circ} + \frac{\sin 147^\circ}{\cos 147^\circ}$ is equal to :
- (1) 0 (2) -1 (3) 1 (4) None of these
8. $P(n) : 1 + 3 + 5 + \dots + (2n - 1) = n^2$ is :
- (1) true for $n > 1$ (2) true for no n
(3) true for all $n \in N$ (4) none of these
9. If eleven members of a committee sit at a round table so that the President and Cashier always sit together, then the number of arrangements is :
- (1) $\underline{10} \times 2$ (2) $\underline{9} \times 2$ (3) $\underline{10}$ (4) None of these
10. In how many ways can 5 keys be put in a ring ?
- (1) $\underline{5}$ (2) $\frac{\underline{5}}{2}$ (3) $\underline{4}$ (4) $\frac{\underline{4}}{2}$

11. A lady gives a dinner party for six guests. The number of ways in which they may be selected from among ten friends, if two of the friends will not attend the party together, is :
 (1) 164 (2) 140 (3) 112 (4) 40
12. If w is an imaginary cube root of unity, then $(1 + w - w^2)$ is equal to :
 (1) $128w$ (2) $-128w$ (3) $-128w^2$ (4) $128w^2$
13. The equation $z\bar{z} + (2 - 3i)z + (2 + 3i)\bar{z} + 4 = 0$ represents a circle of radius :
 (1) 3 (2) 4 (3) 2 (4) 6
14. If the roots of the equation $qx^2 + px + q = 0$ are complex, where p, q are real; then the roots of the equation $x^2 - 4qx + p^2 = 0$ are :
 (1) real and equal (2) imaginary
 (3) real and unequal (4) none of these
15. If $a + b = 8$, then ab is greatest when :
 (1) $a = 3, b = 5$ (2) $a = 4, b = 4$ (3) $a = 6, b = 2$ (4) None of these
16. If the coefficient of 7th and 13th term in the expansion of $(1+x)^n$ are equal, then n is equal to :
 (1) 10 (2) 20 (3) 15 (4) 18
17. If ${}^n C_r$ denotes the number of combinations of n things taken r at a time, then the expression ${}^n C_{r+1} + {}^n C_{r-1} + 2 \times {}^n C_r$ equals :
 (1) ${}^{n+2} C_r$ (2) ${}^{n+2} C_{r+1}$ (3) ${}^{n+1} C_r$ (4) ${}^{n+1} C_{r+1}$
18. The number which should be added to the numbers 2, 14, 62, so that the resulting numbers may be in G. P. is :
 (1) 4 (2) 3 (3) 2 (4) 1
19. If the roots of the equation $x^3 - 12x^2 + 39x - 28 = 0$ are in A. P., then their common difference is :
 (1) ± 2 (2) ± 4 (3) ± 1 (4) ± 3
20. The equation of the straight line joining the origin to the point of intersection of $y - x + 7 = 0$ and $y + 2x - 2 = 0$ is :
 (1) $3x + 4y = 0$ (2) $4x + 3y = 0$ (3) $3x - 4y = 0$ (4) $4x - 3y = 0$

21. The straight line whose sum of the intercepts on the axes is equal to half of the product of the intercepts, passes through the point :
- (1) (2, 2) (2) (1, 1) (3) (4, 4) (4) (3, 3)
22. The equation of a circle with centre (1, 2) and tangent $x + y - 5 = 0$ is :
- (1) $x^2 + y^2 + 2x - 4y + 6 = 0$ (2) $x^2 + y^2 - 2x - 4y + 3 = 0$
 (3) $x^2 + y^2 - 2x + 4y + 8 = 0$ (4) $x^2 + y^2 - 2x - 4y + 8 = 0$
23. The distance between the foci of an ellipse is 16 and the eccentricity is $\frac{1}{2}$. Length of major axis of the ellipse is :
- (1) 8 (2) 16 (3) 32 (4) 64
24. The ratio in which the line joining (2, 4, 5) and (3, 5, -4) is divided by the yz -plane is :
- (1) 4 : -3 (2) 3 : 2 (3) 2 : 3 (4) -2 : 3
25. A plane makes intercepts 3 and 4 respectively on z -axis and x -axis. If plane is parallel to y -axis, then its equation is :
- (1) $3z + 4x = 12$ (2) $3y + 4z = 12$ (3) $3x + 4z = 12$ (4) $3z + 4y = 12$
26. $\lim_{x \rightarrow 0} \frac{\int_0^x y \, dy}{x \tan(\pi + x)}$ is equal to :
- (1) 2 (2) $\frac{1}{2}$ (3) -2 (4) None of these
27. The points of discontinuity of $\tan x$ are :
- (1) $x = n\pi$ (2) $2n\pi$ (3) $(2n+1)\frac{\pi}{2}$ (4) $-2n\pi$
 where $n \in I$
28. If $xy = e^{x-y}$, then $\frac{dy}{dx}$ is equal to :
- (1) $\frac{(x-1)y}{x(1+y)}$ (2) $\frac{(x+1)y}{x(1+y)}$ (3) $\frac{(x-1)y}{x(1-y)}$ (4) None of these
29. If $y = \frac{1 + \sin x - \cos x}{1 + \sin x + \cos x}$, then $\frac{dy}{dx}$ is equal to :
- (1) $\frac{1}{\cos x}$ (2) $\frac{1}{\sin x}$ (3) $\frac{1}{1 - \cos x}$ (4) None of these

30. If $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$, then A^{100} is equal to :
- (1) $2^{100}A$ (2) $100A$ (3) $2^{99}A$ (4) $299A$
31. If A is a square matrix, then $(A + A')$ is :
- (1) unit matrix (2) symmetric matrix
(3) non-singular matrix (4) skew-symmetric matrix
32. If $A = \begin{bmatrix} \alpha & 0 \\ 1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 0 \\ 5 & 1 \end{bmatrix}$, then value of α for which $A^2 = B$ is :
- (1) 1 (2) -1 (3) 4 (4) no real value
33. If $A = \begin{bmatrix} 0 & 3 \\ 2 & 0 \end{bmatrix}$ and $A^{-1} = \lambda(\text{adj } A)$, then λ is equal to :
- (1) $-\frac{1}{6}$ (2) $\frac{1}{6}$ (3) $\frac{1}{3}$ (4) $-\frac{1}{3}$
34. If A is a square matrix such that $AA' = I = A'A$, then $|A|$ is equal to :
- (1) 0 (2) ± 2 (3) ± 1 (4) None of these
35. If w is a complex cube root of unity, then $\begin{vmatrix} 1 & w & -w^2/2 \\ 1 & 1 & 1 \\ 1 & -1 & 0 \end{vmatrix}$ is equal to :
- (1) 1 (2) w (3) 0 (4) w^2
36. If $C = 2 \cos \theta$, then the value of the determinant $\Delta = \begin{vmatrix} C & 1 & 0 \\ 1 & C & 1 \\ 6 & 1 & C \end{vmatrix}$ is :
- (1) $\frac{\sin 4\theta}{\sin \theta}$ (2) $4 \cos^2 \theta (2 \cos \theta - 1)$
(3) $\frac{2 \sin^2 2\theta}{\sin \theta}$ (4) None of these
37. $x + ky - z = 0$, $3x - ky - z = 0$ and $x - 3y + z = 0$ has non-zero solution for k is equal to :
- (1) 0 (2) 1 (3) -1 (4) None of these

38. If $A = \begin{bmatrix} 1 & 2 \\ 3 & -5 \end{bmatrix}$, then A^{-1} is equal to :

- (1) $\begin{bmatrix} \frac{5}{11} & \frac{2}{11} \\ \frac{3}{11} & -\frac{1}{11} \end{bmatrix}$ (2) $\begin{bmatrix} -\frac{5}{11} & \frac{2}{11} \\ \frac{3}{11} & \frac{1}{11} \end{bmatrix}$ (3) $\begin{bmatrix} \frac{5}{11} & -\frac{2}{11} \\ -\frac{3}{11} & -\frac{1}{11} \end{bmatrix}$ (4) None of these

39. If $A = \begin{bmatrix} 1 & \log_a^b \\ \log_b^a & 1 \end{bmatrix}$, then $|A|$ is equal to :

- (1) 0 (2) 1 (3) \log_a^b (4) \log_b^a

40. If $y = \sqrt{\sin x + \sqrt{\sin x + \sqrt{\sin x + \dots \infty}}}$, then $\frac{dy}{dx}$ is equal to :

- (1) $-\frac{\cos x}{2y-1}$ (2) $\frac{\sin x}{1-2y}$ (3) $\frac{\cos x}{2y-1}$ (4) None of these

41. If $f(x) = x + 2$, then $f''[f(x)]$ at $x = 4$ is :

- (1) 8 (2) 1 (3) 4 (4) 5

42. The value of $\frac{d}{dx} \left[\left(\frac{\tan^2 2x - \tan^2 x}{1 - \tan^2 2x \tan^2 x} \right) \cot 3x \right]$ is :

- (1) $\sec x$ (2) $\sec^3 x$ (3) $\sec x \tan x$ (4) $\sec^2 x$

43. If $x = a \sin \theta$, $y = b \cos \theta$, then $\frac{d^2y}{dx^2}$ is equal to :

- (1) $\frac{b}{a^2} \sec \theta$ (2) $-\frac{b}{a^2} \sec^3 \theta$ (3) $\frac{b}{a} \sec^2 \theta$ (4) None of these

44. If $x^m y^n = (x+y)^{m+n}$, then $\frac{dy}{dx}$ is :

- (1) $\frac{y}{x}$ (2) $\frac{x}{y}$ (3) xy (4) None of these

45. Maximum slope of the curve $y = -x^3 + 3x^2 + 9x - 27$ is :

- (1) 0 (2) 16 (3) 12 (4) 32

46. The function x^x is increasing, when :

- (1) $x > \frac{1}{e}$ (2) $x < \frac{1}{e}$ (3) $x < 0$ (4) None of these

47. The rate of change of the surface area of a sphere of radius r , when the radius is increasing at the rate of 2 cm/s is proportional to :

- (1) $\frac{1}{r}$ (2) r^2 (3) r (4) $\frac{1}{r^2}$

48. Angle between the tangents to the curve $y = x^2 - 5x + 6$ at the points (2, 0) and (3, 0) is :

- (1) $\frac{\pi}{3}$ (2) $\frac{\pi}{2}$ (3) $\frac{\pi}{6}$ (4) $\frac{\pi}{4}$

49. $\int \frac{(x+1)^2}{x(x^2+1)} dx$ is equal to :

- (1) $\log x + C$ (2) $2 \tan^{-1} x + C$ (3) $\log \frac{1}{1+x^2} + C$ (4) None of these

50. $\int \frac{x + \sin x}{1 + \cos x} dx$ is equal to :

- (1) $x \tan \frac{x}{2} + C$ (2) $\tan \frac{x}{2} + C$ (3) $\log \cos \frac{x}{2}$ (4) None of these

51. $\int x^x (1 + \log x) dx$ is equal to :

- (1) $x^{2x} + C$ (2) $x^x + C$ (3) $x^x \log x$ (4) None of these

52. $\int \frac{x}{x^2 + 4x + 5} dx$ is equal to :

(1) $\frac{1}{2} \log [x^2 + 4x + 5] - 2 \tan^{-1}(x+2) + C$

(2) $\frac{1}{2} \log [(x+2)^2 - 1] + 2 \tan^{-1}(x+2) + C$

(3) $\frac{1}{2} \log [x^2 + 4x + 5] + C$

(4) None of these

53. $\int \frac{dx}{\sin x \cos x}$ is equal to :

(1) $\log |\sin x| + C$

(2) $\log |\sec x| + C$

(3) $\log |\tan x| + C$

(4) None of these

54. $\int_{-\frac{1}{2}}^{\frac{1}{2}} (\cos x) \left[\log \left(\frac{1-x}{1+x} \right) \right] dx$ is equal to :
- (1) 1 (2) $e^{\frac{1}{2}}$ (3) 0 (4) -1
55. $\int_2^3 \frac{dx}{x^2-x}$ is equal to :
- (1) $\log \left(\frac{2}{3} \right)$ (2) $\log \left(\frac{1}{4} \right)$ (3) $\log \left(\frac{8}{3} \right)$ (4) $\log \left(\frac{4}{3} \right)$
56. The value of $\int_{-\pi}^{\pi} (1-x^2) \sin x \cos^2 x dx$ is :
- (1) π (2) 0 (3) 2π (4) None of these
57. The area bounded by $y = \log x$, x -axis and ordinates $x = 1$, $x = 2$ is :
- (1) $\log \left(\frac{4}{e} \right)$ sq. unit (2) $\log \left(\frac{2}{e} \right)$ sq. unit
 (3) $\log 4$ sq. unit (4) None of these
58. Area bounded by the curves $y = x^2$ and $y^2 = x$ is :
- (1) $\frac{2}{3}$ sq. unit (2) $\frac{1}{3}$ sq. unit (3) $\frac{1}{2}$ sq. unit (4) None of these
59. Area of region satisfying $x \leq 2$, $y \leq |x|$ and $x \geq 0$ is :
- (1) 4 sq. unit (2) 1 sq. unit (3) 2 sq. unit (4) None of these
60. $\int \frac{dx}{x^2+4x+13}$ is equal to :
- (1) $\tan^{-1} \left(\frac{x+2}{3} \right) + C$ (2) $\frac{1}{3} \tan^{-1} \frac{x}{2} + C$
 (3) $\tan^{-1}(x-2) + C$ (4) None of these
61. If $\vec{a} = \hat{i} + \hat{j} + \hat{k}$, $\vec{b} = 2\hat{i} - 4\hat{k}$, $\vec{c} = \hat{i} + \lambda\hat{j} + 3\hat{k}$ are coplanar, then the value of λ is :
- (1) $\frac{5}{3}$ (2) $\frac{3}{5}$ (3) $\frac{5}{2}$ (4) $\frac{7}{3}$

62. If $\vec{a} = 2\hat{i} + \hat{j} + 2\hat{k}$ and $\vec{b} = 5\hat{i} - 3\hat{j} + \hat{k}$, then the projection of \vec{b} on \vec{a} is :
 (1) 6 (2) 5 (3) 4 (4) 3
63. If $\vec{a}, \vec{b}, \vec{c}$ are mutually perpendicular unit vectors, then $|\vec{a} + \vec{b} + \vec{c}|$ is equal to :
 (1) 3 (2) $\sqrt{3}$ (3) 1 (4) 0
64. If $\vec{a} = \hat{i} + \hat{j} + \hat{k}, \vec{b} = \hat{i} + \hat{j}, \vec{c} = \hat{i}$ and $(\vec{a} \times \vec{b}) \times \vec{c} = \lambda \vec{a} + \mu \vec{b}$, then $\lambda + \mu$ is equal to :
 (1) 3 (2) 1 (3) 0 (4) 2
65. A variable plane moves, so that the sum of the reciprocals of its intercepts on the coordinates axes is $\frac{1}{2}$. Then the plane passes through :
 (1) $\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}\right)$ (2) $(-1, 1, 1)$ (3) $(2, 2, 2)$ (4) $(0, 0, 0)$
66. The equation of the plane which bisects the line joining $(2, 3, 4)$ and $(6, 7, 8)$ is :
 (1) $x + y + z - 15 = 0$ (2) $x + y + z + 15 = 0$
 (3) $x - y - z - 15 = 0$ (4) $x - y + z - 15 = 0$
67. The direction ratio of normal to the plane through $(1, 0, 0), (0, 1, 0)$ which makes an angle $\frac{\pi}{4}$ with plane $x + y = 3$ are :
 (1) $1, \sqrt{2}, 1$ (2) $1, 1, \sqrt{2}$ (3) $1, 1, 2$ (4) $\sqrt{2}, 1, 1$
68. A line makes the same angle θ , with each of the x and z axes. If the angle β which it makes with y -axis is such that $\sin^2 \beta = 3 \sin^2 \theta$, then $\cos^2 \theta$ is equal to :
 (1) $\frac{2}{5}$ (2) $\frac{1}{5}$ (3) $\frac{2}{3}$ (4) $\frac{3}{5}$
69. The solution of $\frac{dy}{dx} = 2^{y-x}$ is :
 (1) $2^x + 2^y = C$ (2) $\frac{1}{2^x} - \frac{1}{2^y} = C$ (3) $2^x - 2^y = C$ (4) None of these
70. $y + x^2 = \frac{dy}{dx}$ has the solution :
 (1) $y + x^2 + 2x + 2 = ce^x$ (2) $y + 2x = ce^x$
 (3) $y + 2x + 2 = ce^x$ (4) None of these

71. The solution of $x dy - y dx + x^2 e^x dx = 0$ is :
- (1) $\frac{x}{y} + e^x = C$ (2) $x + e^y = C$ (3) $\frac{y}{x} + e^x = C$ (4) $y + e^x = C$
72. The algebraic sum of deviation of 20 observations measured from 30 is 2. The mean observation is :
- (1) 28.5 (2) 29.6 (3) 30.5 (4) 30.1
73. The standard deviation of 15 items is 6 and if each item is decreased by 1, then standard deviation will be :
- (1) 5 (2) 7 (3) 6 (4) None of these
74. If in a frequency distribution, the mean and median are 21 and 22 respectively, then its mode is approximately :
- (1) 24 (2) 42 (3) 22 (4) 20
75. A coin is tossed 4 times. The probability that at least one head turns up is :
- (1) $\frac{1}{16}$ (2) $\frac{15}{16}$ (3) $\frac{2}{16}$ (4) None of these
76. One card is drawn randomly from a pack of 52 cards, then the probability that it is a king or spade is :
- (1) $\frac{1}{13}$ (2) $\frac{2}{13}$ (3) $\frac{3}{13}$ (4) $\frac{4}{13}$
77. A problem in mathematics is given to three students A, B, C and their respective probability of solving the problem is $\frac{1}{2}$, $\frac{1}{3}$ and $\frac{1}{4}$. Probability that the problem is solved, is :
- (1) $\frac{1}{3}$ (2) $\frac{1}{2}$ (3) $\frac{3}{4}$ (4) $\frac{2}{3}$
78. Five coins whose faces are marked 2, 3 are tossed. The chance of obtaining a total of 12 is :
- (1) $\frac{1}{32}$ (2) $\frac{1}{16}$ (3) $\frac{3}{16}$ (4) $\frac{5}{16}$
79. A card is drawn from a pack of cards. The probability that the card will be a queen or a heart, is :
- (1) $\frac{2}{13}$ (2) $\frac{4}{13}$ (3) $\frac{3}{13}$ (4) None of these

80. If a dice is thrown twice, the probability of occurrence of 4 at least once, is :
 (1) $\frac{11}{36}$ (2) $\frac{7}{12}$ (3) $\frac{35}{36}$ (4) None of these
81. Five horses are in the race. Mr. B selects two of the horses at random and bets on them. The probability that Mr. B selected the winning horse, is :
 (1) $\frac{1}{5}$ (2) $\frac{2}{5}$ (3) $\frac{3}{5}$ (4) None of these
82. If A and B are events such that $P(A \cup B) = \frac{3}{4}$, $P(A \cap B) = \frac{1}{4}$, $P(\bar{A}) = \frac{2}{3}$, then $P(\bar{A} \cap B)$ is :
 (1) $\frac{5}{12}$ (2) $\frac{3}{8}$ (3) $\frac{5}{8}$ (4) $\frac{1}{4}$
83. A coin is tossed three times. The probability of getting head and tail alternatively, is :
 (1) $\frac{1}{8}$ (2) $\frac{1}{2}$ (3) $\frac{1}{4}$ (4) None of these
84. Seven white balls and three black balls are randomly placed in a row. The probability that no two black balls are placed adjacently, equals :
 (1) $\frac{1}{13}$ (2) $\frac{2}{15}$ (3) $\frac{7}{15}$ (4) $\frac{1}{2}$
85. The solution set of the equation $\sin^{-1} x = 2 \tan^{-1} x$ is :
 (1) $\{1, 2\}$ (2) $\{-1, 2\}$ (3) $\{-1, 1, 0\}$ (4) $\left\{1, \frac{1}{2}, 0\right\}$
86. $\tan \left[\frac{1}{2} \sin^{-1} \left(\frac{2a}{1+a^2} \right) + \frac{1}{2} \cos^{-1} \left(\frac{1-a^2}{1+a^2} \right) \right]$ is equal to :
 (1) $\frac{2a}{1+a^2}$ (2) $\frac{2a}{1-a^2}$ (3) $\frac{1-a^2}{1+a^2}$ (4) $\frac{1+a^2}{1-a^2}$
87. $\tan^{-1} \frac{x}{y} - \tan^{-1} \frac{x-y}{x+y}$; ($x > y > 0$) is equal to :
 (1) $\frac{3\pi}{4}$ (2) $-\frac{\pi}{4}$ (3) $-\frac{3\pi}{4}$ (4) $\frac{\pi}{4}$
88. If in a triangle ABC , $\angle A = \tan^{-1} 2$ and $\angle B = \tan^{-1} 3$, then angle C is equal to :
 (1) $\frac{\pi}{4}$ (2) $\frac{3\pi}{4}$ (3) $-\frac{\pi}{4}$ (4) None of these

89. For real numbers x and y , we write $xRy \Leftrightarrow x^2 - y^2 + \sqrt{3}$ is an irrational number. Then the relation R , is :
- (1) Transitive (2) Reflexive (3) Symmetric (4) None of these
90. A function f from the set of natural numbers to integers defined by
- $$f(x) = \begin{cases} \frac{n-1}{2}, & \text{when } n \text{ is odd} \\ -\frac{n}{2}, & \text{when } n \text{ is even} \end{cases} \text{ is :}$$
- (1) one-one but not onto (2) onto but not one-one
(3) one-one and onto both (4) none of these
91. Let R be the relation from $A = \{2, 3, 4, 5\}$ to $B = \{3, 6, 7, 10\}$ defined by 'x divides y', then R^{-1} is equal to
- (1) $\{(6, 2), (3, 3)\}$ (2) $\{(6, 2), (10, 2)\}$
(3) $\{(6, 2), (10, 2), (3, 3), (6, 3), (10, 5)\}$ (4) None of these
92. Which of the following is a singleton set ?
- (1) $\{x : |x| < 1, x \in \mathbb{Z}\}$ (2) $\{x : |x| = 5, x \in \mathbb{Z}\}$
(3) $\{x : x^2 = 1, x \in \mathbb{Z}\}$ (4) $\{x : x^2 + x + 1 = 0, x \in \mathbb{R}\}$
93. If $A = \{(x, y) : y = e^x, x \in \mathbb{R}\}$ and $B = \{(x, y) : y = e^{-x}, x \in \mathbb{R}\}$, then $A \cap B$ is :
- (1) empty set (2) not a set (3) singleton set (4) none of these
94. If $A = \{(x, y) : x^2 + y^2 = 25\}$ and $B = \{(x, y) : x^2 + 9y^2 = 144\}$, then $A \cap B$ contains :
- (1) one point (2) two points (3) three points (4) four points
95. If $z = i \log(2 - \sqrt{3})$, then $\cos z$ is equal to :
- (1) i (2) 2 (3) $3i$ (4) $2i$
96. The expression $\tan^2 \alpha + \cot^2 \alpha$ is :
- (1) ≥ 2 (2) ≤ 2 (3) ≥ -2 (4) None of these
97. A linear programming problem is concerned with finding the following value :
- (1) only maximum value (2) optimal value
(3) only minimum value (4) none of these

98. The linear function $Z = ax + by$, where a, b are constants, which has to be maximized or minimized is called a :
- (1) constraint (2) function of any type
(3) linear objective function (4) none of these
99. A compound statement is a statement which is made up of :
- (1) only one statement (2) any number of statements
(3) two or more statements (4) none of these
100. A compound statement with an 'Or' is false when :
- (1) one component statement is false
(2) none component statement is false
(3) both the component statements are false
(4) none of these

Used for Evaluation for OMR sheets

Total No. of Printed Pages : 13

(DO NOT OPEN THIS QUESTION BOOKLET BEFORE TIME OR UNTIL YOU ARE ASKED TO DO SO)

PG-EE-2016

SUBJECT : Mathematics Hons. Five Year

B

10090

Sr. No.

Time : 1½ Hours

Max. Marks : 100

Total Questions : 100

Roll No. (in figures) _____ (in words) _____

Name _____ Father's Name _____

Mother's Name _____ Date of Examination _____

(Signature of the Candidate)

(Signature of the Invigilator)

CANDIDATES MUST READ THE FOLLOWING INFORMATION/INSTRUCTIONS BEFORE STARTING THE QUESTION PAPER.

- All questions are compulsory.**
- The candidates **must return** the question booklet as well as OMR Answer-Sheet to the Invigilator concerned before leaving the Examination Hall, failing which a case of use of unfair-means/misbehaviour will be registered against him/her, in addition to lodging of an FIR with the police. Further the answer-sheet of such a candidate will not be evaluated.
- In case there is any discrepancy in any question(s) in the Question Booklet, the same may be brought to the notice of the Controller of Examinations in writing **within two hours** after the test is over. No such complaint(s) will be entertained thereafter.
- The candidate **must not** do any rough work or writing in the OMR Answer-Sheet. Rough work, if any, may be done in the question booklet itself. Answers **must not** be ticked in the question booklet.
- There will be no negative marking. Each correct answer will be awarded one full mark. Cutting, erasing, overwriting and more than one answer in OMR Answer-Sheet will be treated as incorrect answer.**
- Use only **Black** or **Blue Ball Point Pen** of good quality in the OMR Answer-Sheet.
- Before answering the questions, the candidates should ensure that they have been supplied correct and complete booklet. Complaints, if any, regarding misprinting etc. will not be entertained 30 minutes after starting of the examination.**

PG-EE-2016/(Maths Hons Five Year)/(B)

SEAL

1. If $f(x) = x + 2$, then $f'[f(x)]$ at $x = 4$ is :
 (1) 8 (2) 1 (3) 4 (4) 5
2. The value of $\frac{d}{dx} \left[\left(\frac{\tan^2 2x - \tan^2 x}{1 - \tan^2 2x \tan^2 x} \right) \cot 3x \right]$ is :
 (1) $\sec x$ (2) $\sec^3 x$ (3) $\sec x \tan x$ (4) $\sec^2 x$
3. If $x = a \sin \theta$, $y = b \cos \theta$, then $\frac{d^2 y}{dx^2}$ is equal to :
 (1) $\frac{b}{a^2} \sec \theta$ (2) $-\frac{b}{a^2} \sec^3 \theta$ (3) $\frac{b}{a} \sec^2 \theta$ (4) None of these
4. If $x^m y^n = (x+y)^{m+n}$, then $\frac{dy}{dx}$ is :
 (1) $\frac{y}{x}$ (2) $\frac{x}{y}$ (3) xy (4) None of these
5. Maximum slope of the curve $y = -x^3 + 3x^2 + 9x - 27$ is :
 (1) 0 (2) 16 (3) 12 (4) 32
6. The function x^x is increasing, when :
 (1) $x > \frac{1}{e}$ (2) $x < \frac{1}{e}$ (3) $x < 0$ (4) None of these
7. The rate of change of the surface area of a sphere of radius r , when the radius is increasing at the rate of 2 cm/s is proportional to :
 (1) $\frac{1}{r}$ (2) r^2 (3) r (4) $\frac{1}{r^2}$
8. Angle between the tangents to the curve $y = x^2 - 5x + 6$ at the points (2, 0) and (3, 0) is :
 (1) $\frac{\pi}{3}$ (2) $\frac{\pi}{2}$ (3) $\frac{\pi}{6}$ (4) $\frac{\pi}{4}$
9. $\int \frac{(x+1)^2}{x(x^2+1)} dx$ is equal to :
 (1) $\log x + C$ (2) $2 \tan^{-1} x + C$ (3) $\log \frac{1}{1+x^2} + C$ (4) None of these

10. $\int \frac{x + \sin x}{1 + \cos x} dx$ is equal to :
- (1) $x \tan \frac{x}{2} + C$ (2) $\tan \frac{x}{2} + C$ (3) $\log \cos \frac{x}{2}$ (4) None of these
11. The straight line whose sum of the intercepts on the axes is equal to half of the product of the intercepts, passes through the point :
- (1) (2, 2) (2) (1, 1) (3) (4, 4) (4) (3, 3)
12. The equation of a circle with centre (1, 2) and tangent $x + y - 5 = 0$ is :
- (1) $x^2 + y^2 + 2x - 4y + 6 = 0$ (2) $x^2 + y^2 - 2x - 4y + 3 = 0$
 (3) $x^2 + y^2 - 2x + 4y + 8 = 0$ (4) $x^2 + y^2 - 2x - 4y + 8 = 0$
13. The distance between the foci of an ellipse is 16 and the eccentricity is $\frac{1}{2}$. Length of major axis of the ellipse is :
- (1) 8 (2) 16 (3) 32 (4) 64
14. The ratio in which the line joining (2, 4, 5) and (3, 5, -4) is divided by the yz -plane is :
- (1) 4 : -3 (2) 3 : 2 (3) 2 : 3 (4) -2 : 3
15. A plane makes intercepts 3 and 4 respectively on z -axis and x -axis. If plane is parallel to y -axis, then its equation is :
- (1) $3z + 4x = 12$ (2) $3y + 4z = 12$ (3) $3x + 4z = 12$ (4) $3z + 4y = 12$
16. $\lim_{x \rightarrow 0} \frac{\int_0^x y dy}{x \tan(\pi + x)}$ is equal to :
- (1) 2 (2) $\frac{1}{2}$ (3) -2 (4) None of these
17. The points of discontinuity of $\tan x$ are :
- (1) $x = n\pi$ (2) $2n\pi$ (3) $(2n+1)\frac{\pi}{2}$ (4) $-2n\pi$
 where $n \in I$
18. If $xy = e^{x-y}$, then $\frac{dy}{dx}$ is equal to :
- (1) $\frac{(x-1)y}{x(1+y)}$ (2) $\frac{(x+1)y}{x(1+y)}$ (3) $\frac{(x-1)y}{x(1-y)}$ (4) None of these

19. If $y = \frac{1 + \sin x - \cos x}{1 + \sin x + \cos x}$, then $\frac{dy}{dx}$ is equal to :
- (1) $\frac{1}{\cos x}$ (2) $\frac{1}{\sin x}$ (3) $\frac{1}{1 - \cos x}$ (4) None of these
20. If $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$, then A^{100} is equal to :
- (1) $2^{100}A$ (2) $100A$ (3) $2^{99}A$ (4) $299A$
21. If two sets A and B are having 99 elements in common, then the number of elements common to each of the sets $A \times B$ and $B \times A$ are :
- (1) 99^2 (2) 18 (3) 2^{99} (4) 100
22. If $f(x) = 1 - \frac{1}{x}$, then the value of $f\left[f\left(\frac{1}{x}\right)\right]$ is :
- (1) $\frac{1}{x} - 1$ (2) $\frac{x}{1-x}$ (3) $\frac{x-1}{x}$ (4) $\frac{x}{x-1}$
23. The function $f(x) = \log\left(x + \sqrt{x^2 + 1}\right)$ is :
- (1) an even function (2) an odd function
(3) a periodic function (4) None of these
24. The value of $\sin A \sin(60^\circ + A) \sin(60^\circ - A)$ is equal to :
- (1) $\sin 3A$ (2) $\sin \frac{3A}{2}$ (3) $\sin \frac{3A}{4}$ (4) $\sin \frac{4A}{3}$
25. If $\cos \theta = \frac{1}{2}\left(x + \frac{1}{x}\right)$, then $\frac{1}{2}\left(x^2 + \frac{1}{x^2}\right)$ is equal to :
- (1) $\cos 2\theta$ (2) $\sin 2\theta$ (3) $\sec 2\theta$ (4) $\tan 2\theta$
26. If $y = \sin^2 \theta + \operatorname{cosec}^2 \theta$, $\theta \neq 0$, then :
- (1) $y > 2$ (2) $y \leq 2$ (3) $y \geq -2$ (4) $y = 0$
27. The value of $\frac{\cos 12^\circ - \sin 12^\circ}{\cos 12^\circ + \sin 12^\circ} + \frac{\sin 147^\circ}{\cos 147^\circ}$ is equal to :
- (1) 0 (2) -1 (3) 1 (4) None of these

28. $P(n) : 1 + 3 + 5 + \dots + (2n - 1) = n^2$ is :
 (1) true for $n > 1$ (2) true for no n
 (3) true for all $n \in \mathbb{N}$ (4) None of these
29. If eleven members of a committee sit at a round table so that the President and Cashier always sit together, then the number of arrangements is :
 (1) $\underline{10} \times 2$ (2) $\underline{9} \times 2$ (3) $\underline{10}$ (4) None of these
30. In how many ways can 5 keys be put in a ring ?
 (1) $\underline{5}$ (2) $\frac{\underline{5}}{2}$ (3) $\underline{4}$ (4) $\frac{\underline{4}}{2}$
31. Let R be the relation from $A = \{2, 3, 4, 5\}$ to $B = \{3, 6, 7, 10\}$ defined by 'x divides y', then R^{-1} is equal to :
 (1) $\{(6, 2), (3, 3)\}$ (2) $\{(6, 2), (10, 2)\}$
 (3) $\{(6, 2), (10, 2), (3, 3), (6, 3), (10, 5)\}$ (4) None of these
32. Which of the following is a singleton set ?
 (1) $\{x : |x| < 1, x \in \mathbb{Z}\}$ (2) $\{x : |x| = 5, x \in \mathbb{Z}\}$
 (3) $\{x : x^2 = 1, x \in \mathbb{Z}\}$ (4) $\{x : x^2 + x + 1 = 0, x \in \mathbb{R}\}$
33. If $A = \{(x, y) : y = e^x, x \in \mathbb{R}\}$ and $B = \{(x, y) : y = e^{-x}, x \in \mathbb{R}\}$, then $A \cap B$ is :
 (1) empty set (2) not a set (3) singleton set (4) none of these
34. If $A = \{(x, y) : x^2 + y^2 = 25\}$ and $B = \{(x, y) : x^2 + 9y^2 = 144\}$, then $A \cap B$ contains :
 (1) one point (2) two points (3) three points (4) four points
35. If $z = i \log(2 - \sqrt{3})$, then $\cos z$ is equal to :
 (1) i (2) 2 (3) $3i$ (4) $2i$
36. The expression $\tan^2 \alpha + \cot^2 \alpha$ is :
 (1) ≥ 2 (2) ≤ 2 (3) ≥ -2 (4) None of these
37. A linear programming problem is concerned with finding the following value :
 (1) only maximum value (2) optimal value
 (3) only minimum value (4) none of these

38. The linear function $Z = ax + by$, where a, b are constants, which has to be maximized or minimized is called a :
- (1) constraint (2) function of any type
(3) linear objective function (4) none of these
39. A compound statement is a statement which is made up of :
- (1) only one statement (2) any number of statements
(3) two or more statements (4) none of these
40. A compound statement with an 'Or' is false when :
- (1) one component statement is false
(2) none component statement is false
(3) both the component statements are false
(4) none of these
41. If $\vec{a} = \hat{i} + \hat{j} + \hat{k}$, $\vec{b} = 2\hat{i} - 4\hat{k}$, $\vec{c} = \hat{i} + \lambda\hat{j} + 3\hat{k}$ are coplanar, then the value of λ is :
- (1) $\frac{5}{3}$ (2) $\frac{3}{5}$ (3) $\frac{5}{2}$ (4) $\frac{7}{3}$
42. If $\vec{a} = 2\hat{i} + \hat{j} + 2\hat{k}$ and $\vec{b} = 5\hat{i} - 3\hat{j} + \hat{k}$, then the projection of \vec{b} on \vec{a} is :
- (1) 6 (2) 5 (3) 4 (4) 3
43. If $\vec{a}, \vec{b}, \vec{c}$ are mutually perpendicular unit vectors, then $|\vec{a} + \vec{b} + \vec{c}|$ is equal to :
- (1) 3 (2) $\sqrt{3}$ (3) 1 (4) 0
44. If $\vec{a} = \hat{i} + \hat{j} + \hat{k}$, $\vec{b} = \hat{i} + \hat{j}$, $\vec{c} = \hat{i}$ and $(\vec{a} \times \vec{b}) \times \vec{c} = \lambda \vec{a} + \mu \vec{b}$, then $\lambda + \mu$ is equal to :
- (1) 3 (2) 1 (3) 0 (4) 2
45. A variable plane moves, so that the sum of the reciprocals of its intercepts on the coordinates axes is $\frac{1}{2}$. Then the plane passes through :
- (1) $\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}\right)$ (2) $(-1, 1, 1)$ (3) $(2, 2, 2)$ (4) $(0, 0, 0)$
46. The equation of the plane which bisects the line joining $(2, 3, 4)$ and $(6, 7, 8)$ is :
- (1) $x + y + z - 15 = 0$ (2) $x + y + z + 15 = 0$
(3) $x - y - z - 15 = 0$ (4) $x - y + z - 15 = 0$

47. The direction ratio of normal to the plane through $(1, 0, 0)$, $(0, 1, 0)$ which makes an angle $\frac{\pi}{4}$ with plane $x + y = 3$ are :
- (1) $1, \sqrt{2}, 1$ (2) $1, 1, \sqrt{2}$ (3) $1, 1, 2$ (4) $\sqrt{2}, 1, 1$
48. A line makes the same angle θ , with each of the x and z axes. If the angle β which it makes with y -axis is such that $\sin^2 \beta = 3 \sin^2 \theta$, then $\cos^2 \theta$ is equal to :
- (1) $\frac{2}{5}$ (2) $\frac{1}{5}$ (3) $\frac{2}{3}$ (4) $\frac{3}{5}$
49. The solution of $\frac{dy}{dx} = 2^{y-x}$ is :
- (1) $2^x + 2^y = C$ (2) $\frac{1}{2^x} - \frac{1}{2^y} = C$ (3) $2^x - 2^y = C$ (4) None of these
50. $y + x^2 = \frac{dy}{dx}$ has the solution :
- (1) $y + x^2 + 2x + 2 = ce^x$ (2) $y + 2x = ce^x$
 (3) $y + 2x + 2 = ce^x$ (4) None of these
51. If A is a square matrix, then $(A + A')$ is :
- (1) unit matrix (2) symmetric matrix
 (3) non-singular matrix (4) skew-symmetric matrix
52. If $A = \begin{bmatrix} \alpha & 0 \\ 1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 0 \\ 5 & 1 \end{bmatrix}$, then value of α for which $A^2 = B$ is :
- (1) 1 (2) -1 (3) 4 (4) no real value
53. If $A = \begin{bmatrix} 0 & 3 \\ 2 & 0 \end{bmatrix}$ and $A^{-1} = \lambda(\text{adj } A)$, then λ is equal to :
- (1) $-\frac{1}{6}$ (2) $\frac{1}{6}$ (3) $\frac{1}{3}$ (4) $-\frac{1}{3}$
54. If A is a square matrix such that $AA' = I = A'A$, then $|A|$ is equal to :
- (1) 0 (2) ± 2 (3) ± 1 (4) None of these

55. If w is a complex cube root of unity, then $\begin{vmatrix} 1 & w & -w^2/2 \\ 1 & 1 & 1 \\ 1 & -1 & 0 \end{vmatrix}$ is equal to :

- (1) 1 (2) w (3) 0 (4) w^2

56. If $C = 2 \cos \theta$, then the value of the determinant $\Delta = \begin{vmatrix} C & 1 & 0 \\ 1 & C & 1 \\ 6 & 1 & C \end{vmatrix}$ is :

- (1) $\frac{\sin 4\theta}{\sin \theta}$ (2) $4 \cos^2 \theta (2 \cos \theta - 1)$
 (3) $\frac{2 \sin^2 2\theta}{\sin \theta}$ (4) None of these

57. $x + ky - z = 0$, $3x - ky - z = 0$ and $x - 3y + z = 0$ has non-zero solution for k is equal to :

- (1) 0 (2) 1 (3) -1 (4) None of these

58. If $A = \begin{bmatrix} 1 & 2 \\ 3 & -5 \end{bmatrix}$, then A^{-1} is equal to :

- (1) $\begin{bmatrix} \frac{5}{11} & \frac{2}{11} \\ \frac{3}{11} & -\frac{1}{11} \end{bmatrix}$ (2) $\begin{bmatrix} -\frac{5}{11} & \frac{2}{11} \\ \frac{3}{11} & \frac{1}{11} \end{bmatrix}$ (3) $\begin{bmatrix} \frac{5}{11} & -\frac{2}{11} \\ -\frac{3}{11} & -\frac{1}{11} \end{bmatrix}$ (4) None of these

59. If $A = \begin{bmatrix} 1 & \log_a^b \\ \log_b^a & 1 \end{bmatrix}$, then $|A|$ is equal to :

- (1) 0 (2) 1 (3) \log_b^a (4) \log_a^b

60. If $y = \sqrt{\sin x + \sqrt{\sin x + \sqrt{\sin x + \dots \infty}}}$, then $\frac{dy}{dx}$ is equal to :

- (1) $-\frac{\cos x}{2y-1}$ (2) $\frac{\sin x}{1-2y}$ (3) $\frac{\cos x}{2y-1}$ (4) None of these

61. The solution of $x dy - y dx + x^2 e^x dx = 0$ is :

- (1) $\frac{x}{y} + e^x = C$ (2) $x + e^y = C$ (3) $\frac{y}{x} + e^x = C$ (4) $y + e^x = C$

62. The algebraic sum of deviation of 20 observations measured from 30 is 2. The mean observation is :
(1) 28.5 (2) 29.6 (3) 30.5 (4) 30.1
63. The standard deviation of 15 items is 6 and if each item is decreased by 1, then standard deviation will be :
(1) 5 (2) 7 (3) 6 (4) None of these
64. If in a frequency distribution, the mean and median are 21 and 22 respectively, then its mode is approximately :
(1) 24 (2) 42 (3) 22 (4) 20
65. A coin is tossed 4 times. The probability that at least one head turns up is :
(1) $\frac{1}{16}$ (2) $\frac{15}{16}$ (3) $\frac{2}{16}$ (4) None of these
66. One card is drawn randomly from a pack of 52 cards, then the probability that it is a king or spade is :
(1) $\frac{1}{13}$ (2) $\frac{2}{13}$ (3) $\frac{3}{13}$ (4) $\frac{4}{13}$
67. A problem in mathematics is given to three students A, B, C and their respective probability of solving the problem is $\frac{1}{2}$, $\frac{1}{3}$ and $\frac{1}{4}$. Probability that the problem is solved, is :
(1) $\frac{1}{3}$ (2) $\frac{1}{2}$ (3) $\frac{3}{4}$ (4) $\frac{2}{3}$
68. Five coins whose faces are marked 2, 3 are tossed. The chance of obtaining a total of 12 is :
(1) $\frac{1}{32}$ (2) $\frac{1}{16}$ (3) $\frac{3}{16}$ (4) $\frac{5}{16}$
69. A card is drawn from a pack of cards. The probability that the card will be a queen or a heart, is :
(1) $\frac{2}{13}$ (2) $\frac{4}{13}$ (3) $\frac{3}{13}$ (4) None of these
70. If a dice is thrown twice, the probability of occurrence of 4 at least once, is :
(1) $\frac{11}{36}$ (2) $\frac{7}{12}$ (3) $\frac{35}{36}$ (4) None of these

71. Five horses are in the race. Mr. B selects two of the horses at random and bets on them. The probability that Mr. B selected the winning horse, is :
- (1) $\frac{1}{5}$ (2) $\frac{2}{5}$ (3) $\frac{3}{5}$ (4) None of these
72. If A and B are events such that $P(A \cup B) = \frac{3}{4}$, $P(A \cap B) = \frac{1}{4}$, $P(\bar{A}) = \frac{2}{3}$, then $P(\bar{A} \cap B)$ is :
- (1) $\frac{5}{12}$ (2) $\frac{3}{8}$ (3) $\frac{5}{8}$ (4) $\frac{1}{4}$
73. A coin is tossed three times. The probability of getting head and tail alternatively, is :
- (1) $\frac{1}{8}$ (2) $\frac{1}{2}$ (3) $\frac{1}{4}$ (4) None of these
74. Seven white balls and three black balls are randomly placed in a row. The probability that no two black balls are placed adjacently, equals :
- (1) $\frac{1}{13}$ (2) $\frac{2}{15}$ (3) $\frac{7}{15}$ (4) $\frac{1}{2}$
75. The solution set of the equation $\sin^{-1} x = 2 \tan^{-1} x$ is :
- (1) $[1, 2]$ (2) $\{-1, 2\}$ (3) $\{-1, 1, 0\}$ (4) $\left\{1, \frac{1}{2}, 0\right\}$
76. $\tan\left[\frac{1}{2}\sin^{-1}\left(\frac{2a}{1+a^2}\right) + \frac{1}{2}\cos^{-1}\left(\frac{1-a^2}{1+a^2}\right)\right]$ is equal to :
- (1) $\frac{2a}{1+a^2}$ (2) $\frac{2a}{1-a^2}$ (3) $\frac{1-a^2}{1+a^2}$ (4) $\frac{1+a^2}{1-a^2}$
77. $\tan^{-1}\frac{x}{y} - \tan^{-1}\frac{x-y}{x+y}$; ($x > y > 0$) is equal to :
- (1) $\frac{3\pi}{4}$ (2) $-\frac{\pi}{4}$ (3) $-\frac{3\pi}{4}$ (4) $\frac{\pi}{4}$
78. If in a triangle ABC , $\angle A = \tan^{-1} 2$ and $\angle B = \tan^{-1} 3$, then angle C is equal to :
- (1) $\frac{\pi}{4}$ (2) $\frac{3\pi}{4}$ (3) $-\frac{\pi}{4}$ (4) None of these

79. For real numbers x and y , we write $xRy \Leftrightarrow x^2 - y^2 + \sqrt{3}$ is an irrational number. Then the relation R , is :
- (1) Transitive (2) Reflexive (3) Symmetric (4) None of these
80. A function f from the set of natural numbers to integers defined by
- $$f(x) = \begin{cases} \frac{n-1}{2}, & \text{when } n \text{ is odd} \\ -\frac{n}{2}, & \text{when } n \text{ is even} \end{cases} \text{ is :}$$
- (1) one-one but not onto (2) onto but not one-one
(3) one-one and onto both (4) none of these
81. A lady gives a dinner party for six guests. The number of ways in which they may be selected from among ten friends, if two of the friends will not attend the party together, is :
- (1) 164 (2) 140 (3) 112 (4) 40
82. If w is an imaginary cube root of unity, then $(1+w-w^2)$ is equal to :
- (1) $128w$ (2) $-128w$ (3) $-128w^2$ (4) $128w^2$
83. The equation $z\bar{z} + (2-3i)z + (2+3i)\bar{z} + 4 = 0$ represents a circle of radius :
- (1) 3 (2) 4 (3) 2 (4) 6
84. If the roots of the equation $qx^2 + px + q = 0$ are complex, where p, q are real; then the roots of the equation $x^2 - 4qx + p^2 = 0$ are :
- (1) real and equal (2) imaginary
(3) real and unequal (4) none of these
85. If $a + b = 8$, then ab is greatest when :
- (1) $a = 3, b = 5$ (2) $a = 4, b = 4$ (3) $a = 6, b = 2$ (4) None of these
86. If the coefficient of 7th and 13th term in the expansion of $(1+x)^n$ are equal, then n is equal to :
- (1) 10 (2) 20 (3) 15 (4) 18

87. If ${}^n C_r$ denotes the number of combinations of n things taken r at a time, then the expression ${}^n C_{r+1} + {}^n C_{r-1} + 2 \times {}^n C_r$ equals :
- (1) ${}^{n+2} C_r$ (2) ${}^{n+2} C_{r+1}$ (3) ${}^{n+1} C_r$ (4) ${}^{n+1} C_{r+1}$
88. The number which should be added to the numbers 2, 14, 62, so that the resulting numbers may be in G. P. is :
- (1) 4 (2) 3 (3) 2 (4) 1
89. If the roots of the equation $x^3 - 12x^2 + 39x - 28 = 0$ are in A. P., then their common difference is :
- (1) ± 2 (2) ± 4 (3) ± 1 (4) ± 3
90. The equation of the straight line joining the origin to the point of intersection of $y - x + 7 = 0$ and $y + 2x - 2 = 0$ is :
- (1) $3x + 4y = 0$ (2) $4x + 3y = 0$ (3) $3x - 4y = 0$ (4) $4x - 3y = 0$
91. $\int x^x (1 + \log x) dx$ is equal to :
- (1) $x^{2x} + C$ (2) $x^x + C$ (3) $x^x \log x$ (4) None of these
92. $\int \frac{x}{x^2 + 4x + 5} dx$ is equal to :
- (1) $\frac{1}{2} \log[x^2 + 4x + 5] - 2 \tan^{-1}(x+2) + C$
- (2) $\frac{1}{2} \log[(x+2)^2 - 1] + 2 \tan^{-1}(x+2) + C$
- (3) $\frac{1}{2} \log[x^2 + 4x + 5] + C$
- (4) None of these
93. $\int \frac{dx}{\sin x \cos x}$ is equal to :
- (1) $\log |\sin x| + C$ (2) $\log |\sec x| + C$
- (3) $\log |\tan x| + C$ (4) None of these

94. $\int_{-\frac{1}{2}}^{\frac{1}{2}} (\cos x) \left[\log \left(\frac{1-x}{1+x} \right) \right] dx$ is equal to :
- (1) 1 (2) $e^{\frac{1}{2}}$ (3) 0 (4) -1
95. $\int_2^3 \frac{dx}{x^2-x}$ is equal to :
- (1) $\log\left(\frac{2}{3}\right)$ (2) $\log\left(\frac{1}{4}\right)$ (3) $\log\left(\frac{8}{3}\right)$ (4) $\log\left(\frac{4}{3}\right)$
96. The value of $\int_{-\pi}^{\pi} (1-x^2) \sin x \cos^2 x dx$ is :
- (1) π (2) 0 (3) 2π (4) None of these
97. The area bounded by $y = \log x$, x -axis and ordinates $x = 1$, $x = 2$ is :
- (1) $\log\left(\frac{4}{e}\right)$ sq. unit (2) $\log\left(\frac{2}{e}\right)$ sq. unit
 (3) $\log 4$ sq. unit (4) None of these
98. Area bounded by the curves $y = x^2$ and $y^2 = x$ is :
- (1) $\frac{2}{3}$ sq. unit (2) $\frac{1}{3}$ sq. unit (3) $\frac{1}{2}$ sq. unit (4) None of these
99. Area of region satisfying $x \leq 2$, $y \leq |x|$ and $x \geq 0$ is :
- (1) 4 sq. unit (2) 1 sq. unit (3) 2 sq. unit (4) None of these
100. $\int \frac{dx}{x^2+4x+13}$ is equal to :
- (1) $\tan^{-1}\left(\frac{x+2}{3}\right) + C$ (2) $\frac{1}{3} \tan^{-1} \frac{x}{2} + C$
 (3) $\tan^{-1}(x-2) + C$ (4) None of these

Used for Evaluation for OMR sheets
M = JSP

Total No. of Printed Pages : 13

(DO NOT OPEN THIS QUESTION BOOKLET BEFORE TIME OR UNTIL YOU ARE ASKED TO DO SO)

PG-EE-2016

SUBJECT : Mathematics Hons. Five Year

C

Sr. No. 10099

Time : 1½ Hours Max. Marks : 100 Total Questions : 100

Roll No. (in figures) _____ (in words) _____

Name _____ Father's Name _____

Mother's Name _____ Date of Examination _____

(Signature of the Candidate)

(Signature of the Invigilator)

CANDIDATES MUST READ THE FOLLOWING INFORMATION/INSTRUCTIONS BEFORE STARTING THE QUESTION PAPER.

1. All questions are compulsory.
2. The candidates **must return** the question booklet as well as OMR Answer-Sheet to the Invigilator concerned before leaving the Examination Hall, failing which a case of use of unfair-means/misbehaviour will be registered against him/her, in addition to lodging of an FIR with the police. Further the answer-sheet of such a candidate will not be evaluated.
3. In case there is any discrepancy in any question(s) in the Question Booklet, the same may be brought to the notice of the Controller of Examinations in writing **within two hours** after the test is over. No such complaint(s) will be entertained thereafter.
4. The candidate **must not** do any rough work or writing in the OMR Answer-Sheet. Rough work, if any, may be done in the question booklet itself. Answers **must not** be ticked in the question booklet.
5. **There will be no negative marking. Each correct answer will be awarded one full mark. Cutting, erasing, overwriting and more than one answer in OMR Answer-Sheet will be treated as incorrect answer.**
6. Use only **Black or Blue Ball Point Pen** of good quality in the OMR Answer-Sheet.
7. **Before answering the questions, the candidates should ensure that they have been supplied correct and complete booklet. Complaints, if any, regarding misprinting etc. will not be entertained 30 minutes after starting of the examination.**

PG-EE-2016/(Maths Hons Five Year)/(C)

SEAL

1. A lady gives a dinner party for six guests. The number of ways in which they may be selected from among ten friends, if two of the friends will not attend the party together, is :
 (1) 164 (2) 140 (3) 112 (4) 40
2. If w is an imaginary cube root of unity, then $(1 + w - w^2)$ is equal to :
 (1) $128w$ (2) $-128w$ (3) $-128w^2$ (4) $128w^2$
3. The equation $z\bar{z} + (2 - 3i)z + (2 + 3i)\bar{z} + 4 = 0$ represents a circle of radius :
 (1) 3 (2) 4 (3) 2 (4) 6
4. If the roots of the equation $qx^2 + px + q = 0$ are complex, where p, q are real; then the roots of the equation $x^2 - 4qx + p^2 = 0$ are :
 (1) real and equal (2) imaginary
 (3) real and unequal (4) none of these
5. If $a + b = 8$, then ab is greatest when :
 (1) $a = 3, b = 5$ (2) $a = 4, b = 4$ (3) $a = 6, b = 2$ (4) None of these
6. If the coefficient of 7th and 13th term in the expansion of $(1+x)^n$ are equal, then n is equal to :
 (1) 10 (2) 20 (3) 15 (4) 18
7. If ${}^n C_r$ denotes the number of combinations of n things taken r at a time, then the expression ${}^n C_{r+1} + {}^n C_{r-1} + 2 \times {}^n C_r$ equals :
 (1) ${}^{n+2} C_r$ (2) ${}^{n+2} C_{r+1}$ (3) ${}^{n+1} C_r$ (4) ${}^{n+1} C_{r+1}$
8. The number which should be added to the numbers 2, 14, 62, so that the resulting numbers may be in G. P. is :
 (1) 4 (2) 3 (3) 2 (4) 1
9. If the roots of the equation $x^3 - 12x^2 + 39x - 28 = 0$ are in A. P., then their common difference is :
 (1) ± 2 (2) ± 4 (3) ± 1 (4) ± 3
10. The equation of the straight line joining the origin to the point of intersection of $y - x + 7 = 0$ and $y + 2x - 2 = 0$ is :
 (1) $3x + 4y = 0$ (2) $4x + 3y = 0$ (3) $3x - 4y = 0$ (4) $4x - 3y = 0$

11. Let R be the relation from $A = \{2, 3, 4, 5\}$ to $B = \{3, 6, 7, 10\}$ defined by 'x divides y', then R^{-1} is equal to :
- (1) $\{(6, 2), (3, 3)\}$ (2) $\{(6, 2), (10, 2)\}$
 (3) $\{(6, 2), (10, 2), (3, 3), (6, 3), (10, 5)\}$ (4) None of these
12. Which of the following is a singleton set ?
- (1) $\{x : |x| < 1, x \in \mathbb{Z}\}$ (2) $\{x : |x| = 5, x \in \mathbb{Z}\}$
 (3) $\{x : x^2 = 1, x \in \mathbb{Z}\}$ (4) $\{x : x^2 + x + 1 = 0, x \in \mathbb{R}\}$
13. If $A = \{(x, y) : y = e^x, x \in \mathbb{R}\}$ and $B = \{(x, y) : y = e^{-x}, x \in \mathbb{R}\}$, then $A \cap B$ is :
- (1) empty set (2) not a set (3) singleton set (4) none of these
14. If $A = \{(x, y) : x^2 + y^2 = 25\}$ and $B = \{(x, y) : x^2 + 9y^2 = 144\}$, then $A \cap B$ contains :
- (1) one point (2) two points (3) three points (4) four points
15. If $z = i \log(2 - \sqrt{3})$, then $\cos z$ is equal to :
- (1) i (2) 2 (3) $3i$ (4) $2i$
16. The expression $\tan^2 \alpha + \cot^2 \alpha$ is :
- (1) ≥ 2 (2) ≤ 2 (3) ≥ -2 (4) None of these
17. A linear programming problem is concerned with finding the following value :
- (1) only maximum value (2) optimal value
 (3) only minimum value (4) none of these
18. The linear function $Z = ax + by$, where a, b are constants, which has to be maximized or minimized is called a :
- (1) constraint (2) function of any type
 (3) linear objective function (4) none of these
19. A compound statement is a statement which is made up of :
- (1) only one statement (2) any number of statements
 (3) two or more statements (4) none of these
20. A compound statement with an 'Or' is false when :
- (1) one component statement is false
 (2) none component statement is false
 (3) both the component statements are false
 (4) none of these

21. The solution of $x dy - y dx + x^2 e^x dx = 0$ is :
- (1) $\frac{x}{y} + e^x = C$ (2) $x + e^y = C$ (3) $\frac{y}{x} + e^x = C$ (4) $y + e^x = C$
22. The algebraic sum of deviation of 20 observations measured from 30 is 2. The mean observation is :
- (1) 28.5 (2) 29.6 (3) 30.5 (4) 30.1
23. The standard deviation of 15 items is 6 and if each item is decreased by 1, then standard deviation will be :
- (1) 5 (2) 7 (3) 6 (4) None of these
24. If in a frequency distribution, the mean and median are 21 and 22 respectively, then its mode is approximately :
- (1) 24 (2) 42 (3) 22 (4) 20
25. A coin is tossed 4 times. The probability that at least one head turns up is :
- (1) $\frac{1}{16}$ (2) $\frac{15}{16}$ (3) $\frac{2}{16}$ (4) None of these
26. One card is drawn randomly from a pack of 52 cards, then the probability that it is a king or spade is :
- (1) $\frac{1}{13}$ (2) $\frac{2}{13}$ (3) $\frac{3}{13}$ (4) $\frac{4}{13}$
27. A problem in mathematics is given to three students A, B, C and their respective probability of solving the problem is $\frac{1}{2}$, $\frac{1}{3}$ and $\frac{1}{4}$. Probability that the problem is solved, is :
- (1) $\frac{1}{3}$ (2) $\frac{1}{2}$ (3) $\frac{3}{4}$ (4) $\frac{2}{3}$
28. Five coins whose faces are marked 2, 3 are tossed. The chance of obtaining a total of 12 is :
- (1) $\frac{1}{32}$ (2) $\frac{1}{16}$ (3) $\frac{3}{16}$ (4) $\frac{5}{16}$
29. A card is drawn from a pack of cards. The probability that the card will be a queen or a heart, is :
- (1) $\frac{2}{13}$ (2) $\frac{4}{13}$ (3) $\frac{3}{13}$ (4) None of these

30. If a dice is thrown twice, the probability of occurrence of 4 at least once, is :

- (1) $\frac{11}{36}$ (2) $\frac{7}{12}$ (3) $\frac{35}{36}$ (4) None of these

31. $\int x^x(1 + \log x) dx$ is equal to :

- (1) $x^{2x} + C$ (2) $x^x + C$ (3) $x^x \log x$ (4) None of these

32. $\int \frac{x}{x^2 + 4x + 5} dx$ is equal to :

(1) $\frac{1}{2} \log[x^2 + 4x + 5] - 2 \tan^{-1}(x + 2) + C$

(2) $\frac{1}{2} \log[(x + 2)^2 - 1] + 2 \tan^{-1}(x + 2) + C$

(3) $\frac{1}{2} \log[x^2 + 4x + 5] + C$

(4) None of these

33. $\int \frac{dx}{\sin x \cos x}$ is equal to :

(1) $\log |\sin x| + C$

(2) $\log |\sec x| + C$

(3) $\log |\tan x| + C$

(4) None of these

34. $\int_{-\frac{1}{2}}^{\frac{1}{2}} (\cos x) \left[\log \left(\frac{1-x}{1+x} \right) \right] dx$ is equal to :

(1) 1

(2) $e^{\frac{1}{2}}$

(3) 0

(4) -1

35. $\int_2^3 \frac{dx}{x^2 - x}$ is equal to :

(1) $\log \left(\frac{2}{3} \right)$

(2) $\log \left(\frac{1}{4} \right)$

(3) $\log \left(\frac{8}{3} \right)$

(4) $\log \left(\frac{4}{3} \right)$

36. The value of $\int_{-\pi}^{\pi} (1 - x^2) \sin x \cos^2 x dx$ is :

(1) π

(2) 0

(3) 2π

(4) None of these

37. The area bounded by $y = \log x$, x -axis and ordinates $x = 1$, $x = 2$ is :

- (1) $\log\left(\frac{4}{e}\right)$ sq. unit (2) $\log\left(\frac{2}{e}\right)$ sq. unit
(3) $\log 4$ sq. unit (4) None of these

38. Area bounded by the curves $y = x^2$ and $y^2 = x$ is :

- (1) $\frac{2}{3}$ sq. unit (2) $\frac{1}{3}$ sq. unit (3) $\frac{1}{2}$ sq. unit (4) None of these

39. Area of region satisfying $x \leq 2$, $y \leq |x|$ and $x \geq 0$ is :

- (1) 4 sq. unit (2) 1 sq. unit (3) 2 sq. unit (4) None of these

40. $\int \frac{dx}{x^2 + 4x + 13}$ is equal to :

- (1) $\tan^{-1}\left(\frac{x+2}{3}\right) + C$ (2) $\frac{1}{3} \tan^{-1} \frac{x}{2} + C$
(3) $\tan^{-1}(x-2) + C$ (4) None of these

41. If A is a square matrix, then $(A + A')$ is :

- (1) unit matrix (2) symmetric matrix
(3) non-singular matrix (4) skew-symmetric matrix

42. If $A = \begin{bmatrix} \alpha & 0 \\ 1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 0 \\ 5 & 1 \end{bmatrix}$, then value of α for which $A^2 = B$ is :

- (1) 1 (2) -1 (3) 4 (4) no real value

43. If $A = \begin{bmatrix} 0 & 3 \\ 2 & 0 \end{bmatrix}$ and $A^{-1} = \lambda(\text{adj } A)$, then λ is equal to :

- (1) $-\frac{1}{6}$ (2) $\frac{1}{6}$ (3) $\frac{1}{3}$ (4) $-\frac{1}{3}$

44. If A is a square matrix such that $AA' = I = A'A$, then $|A|$ is equal to :

- (1) 0 (2) ± 2 (3) ± 1 (4) None of these

45. If w is a complex cube root of unity, then $\begin{vmatrix} 1 & w & -w^2/2 \\ 1 & 1 & 1 \\ 1 & -1 & 0 \end{vmatrix}$ is equal to :

- (1) 1 (2) w (3) 0 (4) w^2

46. If $C = 2 \cos \theta$, then the value of the determinant $\Delta = \begin{vmatrix} C & 1 & 0 \\ 1 & C & 1 \\ 6 & 1 & C \end{vmatrix}$ is :

- (1) $\frac{\sin 4\theta}{\sin \theta}$ (2) $4 \cos^2 \theta (2 \cos \theta - 1)$
 (3) $\frac{2 \sin^2 2\theta}{\sin \theta}$ (4) None of these

47. $x + ky - z = 0$, $3x - ky - z = 0$ and $x - 3y + z = 0$ has non-zero solution for k is equal to :

- (1) 0 (2) 1 (3) -1 (4) None of these

48. If $A = \begin{bmatrix} 1 & 2 \\ 3 & -5 \end{bmatrix}$, then A^{-1} is equal to :

- (1) $\begin{bmatrix} \frac{5}{11} & \frac{2}{11} \\ \frac{3}{11} & -\frac{1}{11} \end{bmatrix}$ (2) $\begin{bmatrix} -\frac{5}{11} & \frac{2}{11} \\ \frac{3}{11} & \frac{1}{11} \end{bmatrix}$ (3) $\begin{bmatrix} \frac{5}{11} & -\frac{2}{11} \\ -\frac{3}{11} & -\frac{1}{11} \end{bmatrix}$ (4) None of these

49. If $A = \begin{bmatrix} 1 & \log_b^a \\ \log_a^b & 1 \end{bmatrix}$, then $|A|$ is equal to :

- (1) 0 (2) 1 (3) \log_b^a (4) \log_a^b

50. If $y = \sqrt{\sin x + \sqrt{\sin x + \sqrt{\sin x + \dots \infty}}}$, then $\frac{dy}{dx}$ is equal to :

- (1) $-\frac{\cos x}{2y-1}$ (2) $\frac{\sin x}{1-2y}$ (3) $\frac{\cos x}{2y-1}$ (4) None of these

51. The straight line whose sum of the intercepts on the axes is equal to half of the product of the intercepts, passes through the point :

- (1) (2, 2) (2) (1, 1) (3) (4, 4) (4) (3, 3)

52. The equation of a circle with centre (1, 2) and tangent $x + y - 5 = 0$ is :

- (1) $x^2 + y^2 + 2x - 4y + 6 = 0$ (2) $x^2 + y^2 - 2x - 4y + 3 = 0$
 (3) $x^2 + y^2 - 2x + 4y + 8 = 0$ (4) $x^2 + y^2 - 2x - 4y + 8 = 0$

53. The distance between the foci of an ellipse is 16 and the eccentricity is $\frac{1}{2}$. Length of major axis of the ellipse is :

- (1) 8 (2) 16 (3) 32 (4) 64

54. The ratio in which the line joining (2, 4, 5) and (3, 5, -4) is divided by the yz -plane is :

- (1) 4 : -3 (2) 3 : 2 (3) 2 : 3 (4) -2 : 3

55. A plane makes intercepts 3 and 4 respectively on z -axis and x -axis. If plane is parallel to y -axis, then its equation is :

- (1) $3z + 4x = 12$ (2) $3y + 4z = 12$ (3) $3x + 4z = 12$ (4) $3z + 4y = 12$

56. $\lim_{x \rightarrow 0} \frac{\int_0^x y \, dy}{x \tan(\pi + x)}$ is equal to :

- (1) 2 (2) $\frac{1}{2}$ (3) -2 (4) None of these

57. The points of discontinuity of $\tan x$ are :

- (1) $x = n\pi$ (2) $2n\pi$ (3) $(2n+1)\frac{\pi}{2}$ (4) $-2n\pi$

where $n \in I$

58. If $xy = e^{x-y}$, then $\frac{dy}{dx}$ is equal to :

- (1) $\frac{(x-1)y}{x(1+y)}$ (2) $\frac{(x+1)y}{x(1+y)}$ (3) $\frac{(x-1)y}{x(1-y)}$ (4) None of these

59. If $y = \frac{1 + \sin x - \cos x}{1 + \sin x + \cos x}$, then $\frac{dy}{dx}$ is equal to :

- (1) $\frac{1}{\cos x}$ (2) $\frac{1}{\sin x}$ (3) $\frac{1}{1 - \cos x}$ (4) None of these

60. If $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$, then A^{100} is equal to :
- (1) $2^{100}A$ (2) $100A$ (3) $2^{99}A$ (4) $299A$
61. If $f(x) = x + 2$, then $f'[f(x)]$ at $x = 4$ is :
- (1) 8 (2) 1 (3) 4 (4) 5
62. The value of $\frac{d}{dx} \left[\left(\frac{\tan^2 2x - \tan^2 x}{1 - \tan^2 2x \tan^2 x} \right) \cot 3x \right]$ is :
- (1) $\sec x$ (2) $\sec^3 x$ (3) $\sec x \tan x$ (4) $\sec^2 x$
63. If $x = a \sin \theta$, $y = b \cos \theta$, then $\frac{d^2y}{dx^2}$ is equal to :
- (1) $\frac{b}{a^2} \sec \theta$ (2) $-\frac{b}{a^2} \sec^3 \theta$ (3) $\frac{b}{a} \sec^2 \theta$ (4) None of these
64. If $x^m y^n = (x + y)^{m+n}$, then $\frac{dy}{dx}$ is :
- (1) $\frac{y}{x}$ (2) $\frac{x}{y}$ (3) xy (4) None of these
65. Maximum slope of the curve $y = -x^3 + 3x^2 + 9x - 27$ is :
- (1) 0 (2) 16 (3) 12 (4) 32
66. The function x^x is increasing, when :
- (1) $x > \frac{1}{e}$ (2) $x < \frac{1}{e}$ (3) $x < 0$ (4) None of these
67. The rate of change of the surface area of a sphere of radius r , when the radius is increasing at the rate of 2 cm/s is proportional to :
- (1) $\frac{1}{r}$ (2) r^2 (3) r (4) $\frac{1}{r^2}$
68. Angle between the tangents to the curve $y = x^2 - 5x + 6$ at the points (2, 0) and (3, 0) is :
- (1) $\frac{\pi}{3}$ (2) $\frac{\pi}{2}$ (3) $\frac{\pi}{6}$ (4) $\frac{\pi}{4}$

69. $\int \frac{(x+1)^2}{x(x^2+1)} dx$ is equal to :
 (1) $\log x + C$ (2) $2 \tan^{-1} x + C$ (3) $\log \frac{1}{1+x^2} + C$ (4) None of these
70. $\int \frac{x + \sin x}{1 + \cos x} dx$ is equal to :
 (1) $x \tan \frac{x}{2} + C$ (2) $\tan \frac{x}{2} + C$ (3) $\log \cos \frac{x}{2}$ (4) None of these
71. If $\vec{a} = \hat{i} + \hat{j} + \hat{k}$, $\vec{b} = 2\hat{i} - 4\hat{k}$, $\vec{c} = \hat{i} + \lambda\hat{j} + 3\hat{k}$ are coplanar, then the value of λ is :
 (1) $\frac{5}{3}$ (2) $\frac{3}{5}$ (3) $\frac{5}{2}$ (4) $\frac{7}{3}$
72. If $\vec{a} = 2\hat{i} + \hat{j} + 2\hat{k}$ and $\vec{b} = 5\hat{i} - 3\hat{j} + \hat{k}$, then the projection of \vec{b} on \vec{a} is :
 (1) 6 (2) 5 (3) 4 (4) 3
73. If \vec{a} , \vec{b} , \vec{c} are mutually perpendicular unit vectors, then $|\vec{a} + \vec{b} + \vec{c}|$ is equal to :
 (1) 3 (2) $\sqrt{3}$ (3) 1 (4) 0
74. If $\vec{a} = \hat{i} + \hat{j} + \hat{k}$, $\vec{b} = \hat{i} + \hat{j}$, $\vec{c} = \hat{i}$ and $(\vec{a} \times \vec{b}) \times \vec{c} = \lambda \vec{a} + \mu \vec{b}$, then $\lambda + \mu$ is equal to :
 (1) 3 (2) 1 (3) 0 (4) 2
75. A variable plane moves, so that the sum of the reciprocals of its intercepts on the coordinates axes is $\frac{1}{2}$. Then the plane passes through :
 (1) $(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$ (2) $(-1, 1, 1)$ (3) $(2, 2, 2)$ (4) $(0, 0, 0)$
76. The equation of the plane which bisects the line joining $(2, 3, 4)$ and $(6, 7, 8)$ is :
 (1) $x + y + z - 15 = 0$ (2) $x + y + z + 15 = 0$
 (3) $x - y - z - 15 = 0$ (4) $x - y + z - 15 = 0$
77. The direction ratio of normal to the plane through $(1, 0, 0)$, $(0, 1, 0)$ which makes an angle $\frac{\pi}{4}$ with plane $x + y = 3$ are :
 (1) $1, \sqrt{2}, 1$ (2) $1, 1, \sqrt{2}$ (3) $1, 1, 2$ (4) $\sqrt{2}, 1, 1$

78. A line makes the same angle θ , with each of the x and z axes. If the angle β which it makes with y -axis is such that $\sin^2 \beta = 3 \sin^2 \theta$, then $\cos^2 \theta$ is equal to :

- (1) $\frac{2}{5}$ (2) $\frac{1}{5}$ (3) $\frac{2}{3}$ (4) $\frac{3}{5}$

79. The solution of $\frac{dy}{dx} = 2^{y-x}$ is :

- (1) $2^x + 2^y = C$ (2) $\frac{1}{2^x} - \frac{1}{2^y} = C$ (3) $2^x - 2^y = C$ (4) None of these

80. $y + x^2 = \frac{dy}{dx}$ has the solution :

- (1) $y + x^2 + 2x + 2 = ce^x$ (2) $y + 2x = ce^x$
 (3) $y + 2x + 2 = ce^x$ (4) None of these

81. If two sets A and B are having 99 elements in common, then the number of elements common to each of the sets $A \times B$ and $B \times A$ are :

- (1) 99^2 (2) 18 (3) 2^{99} (4) 100

82. If $f(x) = 1 - \frac{1}{x}$, then the value of $f\left[f\left(\frac{1}{x}\right)\right]$ is :

- (1) $\frac{1}{x} - 1$ (2) $\frac{x}{1-x}$ (3) $\frac{x-1}{x}$ (4) $\frac{x}{x-1}$

83. The function $f(x) = \log(x + \sqrt{x^2 + 1})$ is :

- (1) an even function (2) an odd function
 (3) a periodic function (4) none of these

84. The value of $\sin A \sin(60^\circ + A) \sin(60^\circ - A)$ is equal to :

- (1) $\sin 3A$ (2) $\sin \frac{3A}{2}$ (3) $\sin \frac{3A}{4}$ (4) $\sin \frac{4A}{3}$

85. If $\cos \theta = \frac{1}{2}\left(x + \frac{1}{x}\right)$, then $\frac{1}{2}\left(x^2 + \frac{1}{x^2}\right)$ is equal to :

- (1) $\cos 2\theta$ (2) $\sin 2\theta$ (3) $\sec 2\theta$ (4) $\tan 2\theta$

86. If $y = \sin^2 \theta + \operatorname{cosec}^2 \theta$, $\theta \neq 0$, then :

- (1) $y > 2$ (2) $y \leq 2$ (3) $y \geq -2$ (4) $y = 0$

87. The value of $\frac{\cos 12^\circ - \sin 12^\circ}{\cos 12^\circ + \sin 12^\circ} + \frac{\sin 147^\circ}{\cos 147^\circ}$ is equal to :
- (1) 0 (2) -1 (3) 1 (4) None of these
88. $P(n) : 1 + 3 + 5 + \dots + (2n - 1) = n^2$ is :
- (1) true for $n > 1$ (2) true for no n
 (3) true for all $n \in N$ (4) none of these
89. If eleven members of a committee sit at a round table so that the President and Cashier always sit together, then the number of arrangements is :
- (1) $\underline{10} \times 2$ (2) $\underline{9} \times 2$ (3) $\underline{10}$ (4) None of these
90. In how many ways can 5 keys be put in a ring ?
- (1) $\underline{5}$ (2) $\frac{\underline{5}}{2}$ (3) $\underline{4}$ (4) $\frac{\underline{4}}{2}$
91. Five horses are in the race. Mr. B selects two of the horses at random and bets on them. The probability that Mr. B selected the winning horse, is :
- (1) $\frac{1}{5}$ (2) $\frac{2}{5}$ (3) $\frac{3}{5}$ (4) None of these
92. If A and B are events such that $P(A \cup B) = \frac{3}{4}$, $P(A \cap B) = \frac{1}{4}$, $P(\bar{A}) = \frac{2}{3}$, then $P(\bar{A} \cap B)$ is :
- (1) $\frac{5}{12}$ (2) $\frac{3}{8}$ (3) $\frac{5}{8}$ (4) $\frac{1}{4}$
93. A coin is tossed three times. The probability of getting head and tail alternatively, is :
- (1) $\frac{1}{8}$ (2) $\frac{1}{2}$ (3) $\frac{1}{4}$ (4) None of these
94. Seven white balls and three black balls are randomly placed in a row. The probability that no two black balls are placed adjacently, equals :
- (1) $\frac{1}{13}$ (2) $\frac{2}{15}$ (3) $\frac{7}{15}$ (4) $\frac{1}{2}$
95. The solution set of the equation $\sin^{-1} x = 2 \tan^{-1} x$ is :
- (1) $\{1, 2\}$ (2) $\{-1, 2\}$ (3) $\{-1, 1, 0\}$ (4) $\left\{1, \frac{1}{2}, 0\right\}$

96. $\tan\left[\frac{1}{2}\sin^{-1}\left(\frac{2a}{1+a^2}\right) + \frac{1}{2}\cos^{-1}\left(\frac{1-a^2}{1+a^2}\right)\right]$ is equal to :

- (1) $\frac{2a}{1+a^2}$ (2) $\frac{2a}{1-a^2}$ (3) $\frac{1-a^2}{1+a^2}$ (4) $\frac{1+a^2}{1-a^2}$

97. $\tan^{-1}\frac{x}{y} - \tan^{-1}\frac{x-y}{x+y}$; ($x > y > 0$) is equal to :

- (1) $\frac{3\pi}{4}$ (2) $-\frac{\pi}{4}$ (3) $-\frac{3\pi}{4}$ (4) $\frac{\pi}{4}$

98. If in a triangle ABC , $\angle A = \tan^{-1}2$ and $\angle B = \tan^{-1}3$, then angle C is equal to :

- (1) $\frac{\pi}{4}$ (2) $\frac{3\pi}{4}$ (3) $-\frac{\pi}{4}$ (4) None of these

99. For real numbers x and y , we write $xRy \Leftrightarrow x^2 - y^2 + \sqrt{3}$ is an irrational number. Then the relation R , is :

- (1) Transitive (2) Reflexive (3) Symmetric (4) None of these

100. A function f from the set of natural numbers to integers defined by

$$f(x) = \begin{cases} \frac{n-1}{2}, & \text{when } n \text{ is odd} \\ -\frac{n}{2}, & \text{when } n \text{ is even} \end{cases} \text{ is :}$$

- (1) one-one but not onto (2) onto but not one-one
(3) one-one and onto both (4) none of these

Used for Evaluation of OMR Sheets
for Jsgl Merit

Total No. of Printed Pages : 13
(DO NOT OPEN THIS QUESTION BOOKLET BEFORE TIME OR UNTIL YOU ARE ASKED TO DO SO)

PG-EE-2016

SUBJECT : Mathematics Hons. Five Year

D

10108

Sr. No.

Time : 1¼ Hours

Max. Marks : 100

Total Questions : 100

Roll No. (in figures) _____ (in words) _____

Name _____ Father's Name _____

Mother's Name _____ Date of Examination _____

(Signature of the Candidate)

(Signature of the Invigilator)

CANDIDATES MUST READ THE FOLLOWING INFORMATION/INSTRUCTIONS BEFORE STARTING THE QUESTION PAPER.

- 1. All questions are compulsory.**
- 2. The candidates *must return* the question booklet as well as OMR Answer-Sheet to the Invigilator concerned before leaving the Examination Hall, failing which a case of use of unfair-means/misbehaviour will be registered against him/her, in addition to lodging of an FIR with the police. Further the answer-sheet of such a candidate will not be evaluated.**
- 3. In case there is any discrepancy in any question(s) in the Question Booklet, the same may be brought to the notice of the Controller of Examinations in writing **within two hours** after the test is over. No such complaint(s) will be entertained thereafter.**
- 4. The candidate *must not* do any rough work or writing in the OMR Answer-Sheet. Rough work, if any, may be done in the question booklet itself. Answers **must not** be ticked in the question booklet.**
- 5. *There will be no negative marking. Each correct answer will be awarded one full mark. Cutting, erasing, overwriting and more than one answer in OMR Answer-Sheet will be treated as incorrect answer.***
- 6. Use only **Black or Blue Ball Point Pen** of good quality in the OMR Answer-Sheet.**
- 7. *Before answering the questions, the candidates should ensure that they have been supplied correct and complete booklet. Complaints, if any, regarding misprinting etc. will not be entertained 30 minutes after starting of the examination.***

PG-EE-2016/(Maths Hons Five Year)/(D)

1. The solution of $x dy - y dx + x^2 e^x dx = 0$ is :

(1) $\frac{x}{y} + e^x = C$ (2) $x + e^y = C$ (3) $\frac{y}{x} + e^x = C$ (4) $y + e^x = C$

2. The algebraic sum of deviation of 20 observations measured from 30 is 2. The mean observation is :

(1) 28.5 (2) 29.6 (3) 30.5 (4) 30.1

3. The standard deviation of 15 items is 6 and if each item is decreased by 1, then standard deviation will be :

(1) 5 (2) 7 (3) 6 (4) None of these

4. If in a frequency distribution, the mean and median are 21 and 22 respectively, then its mode is approximately :

(1) 24 (2) 42 (3) 22 (4) 20

5. A coin is tossed 4 times. The probability that at least one head turns up is :

(1) $\frac{1}{16}$ (2) $\frac{15}{16}$ (3) $\frac{2}{16}$ (4) None of these

6. One card is drawn randomly from a pack of 52 cards, then the probability that it is a king or spade is :

(1) $\frac{1}{13}$ (2) $\frac{2}{13}$ (3) $\frac{3}{13}$ (4) $\frac{4}{13}$

7. A problem in mathematics is given to three students A, B, C and their respective probability of solving the problem is $\frac{1}{2}$, $\frac{1}{3}$ and $\frac{1}{4}$. Probability that the problem is solved, is :

(1) $\frac{1}{3}$ (2) $\frac{1}{2}$ (3) $\frac{3}{4}$ (4) $\frac{2}{3}$

8. Five coins whose faces are marked 2, 3 are tossed. The chance of obtaining a total of 12 is :

(1) $\frac{1}{32}$ (2) $\frac{1}{16}$ (3) $\frac{3}{16}$ (4) $\frac{5}{16}$

9. A card is drawn from a pack of cards. The probability that the card will be a queen or a heart, is :

(1) $\frac{2}{13}$ (2) $\frac{4}{13}$ (3) $\frac{3}{13}$ (4) None of these

10. If a dice is thrown twice, the probability of occurrence of 4 at least once, is :
 (1) $\frac{11}{36}$ (2) $\frac{7}{12}$ (3) $\frac{35}{36}$ (4) None of these
11. $\int x^x(1 + \log x) dx$ is equal to :
 (1) $x^{2x} + C$ (2) $x^x + C$ (3) $x^x \log x$ (4) None of these
12. $\int \frac{x}{x^2 + 4x + 5} dx$ is equal to :
 (1) $\frac{1}{2} \log[x^2 + 4x + 5] - 2 \tan^{-1}(x+2) + C$
 (2) $\frac{1}{2} \log[(x+2)^2 - 1] + 2 \tan^{-1}(x+2) + C$
 (3) $\frac{1}{2} \log[x^2 + 4x + 5] + C$
 (4) None of these
13. $\int \frac{dx}{\sin x \cos x}$ is equal to :
 (1) $\log |\sin x| + C$ (2) $\log |\sec x| + C$
 (3) $\log |\tan x| + C$ (4) None of these
14. $\int_{-\frac{1}{2}}^{\frac{1}{2}} (\cos x) \left[\log \left(\frac{1-x}{1+x} \right) \right] dx$ is equal to :
 (1) 1 (2) $e^{\frac{1}{2}}$ (3) 0 (4) -1
15. $\int_2^3 \frac{dx}{x^2 - x}$ is equal to :
 (1) $\log \left(\frac{2}{3} \right)$ (2) $\log \left(\frac{1}{4} \right)$ (3) $\log \left(\frac{8}{3} \right)$ (4) $\log \left(\frac{4}{3} \right)$
16. The value of $\int_{-\pi}^{\pi} (1-x^2) \sin x \cos^2 x dx$ is :
 (1) π (2) 0 (3) 2π (4) None of these

17. The area bounded by $y = \log x$, x -axis and ordinates $x = 1$, $x = 2$ is :
- (1) $\log\left(\frac{4}{e}\right)$ sq. unit (2) $\log\left(\frac{2}{e}\right)$ sq. unit
 (3) $\log 4$ sq. unit (4) None of these
18. Area bounded by the curves $y = x^2$ and $y^2 = x$ is :
- (1) $\frac{2}{3}$ sq. unit (2) $\frac{1}{3}$ sq. unit (3) $\frac{1}{2}$ sq. unit (4) None of these
19. Area of region satisfying $x \leq 2$, $y \leq |x|$ and $x \geq 0$ is :
- (1) 4 sq. unit (2) 1 sq. unit (3) 2 sq. unit (4) None of these
20. $\int \frac{dx}{x^2 + 4x + 13}$ is equal to :
- (1) $\tan^{-1}\left(\frac{x+2}{3}\right) + C$ (2) $\frac{1}{3} \tan^{-1} \frac{x}{2} + C$
 (3) $\tan^{-1}(x-2) + C$ (4) None of these
21. If A is a square matrix, then $(A + A')$ is :
- (1) unit matrix (2) symmetric matrix
 (3) non-singular matrix (4) skew-symmetric matrix
22. If $A = \begin{bmatrix} \alpha & 0 \\ 1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 0 \\ 5 & 1 \end{bmatrix}$, then value of α for which $A^2 = B$ is :
- (1) 1 (2) -1 (3) 4 (4) no real value
23. If $A = \begin{bmatrix} 0 & 3 \\ 2 & 0 \end{bmatrix}$ and $A^{-1} = \lambda(\text{adj } A)$, then λ is equal to :
- (1) $-\frac{1}{6}$ (2) $\frac{1}{6}$ (3) $\frac{1}{3}$ (4) $-\frac{1}{3}$
24. If A is a square matrix such that $AA' = I = A'A$, then $|A|$ is equal to :
- (1) 0 (2) ± 2 (3) ± 1 (4) None of these
25. If w is a complex cube root of unity, then $\begin{vmatrix} 1 & w & -w^2/2 \\ 1 & 1 & 1 \\ 1 & -1 & 0 \end{vmatrix}$ is equal to :
- (1) 1 (2) w (3) 0 (4) w^2

26. If $C = 2 \cos \theta$, then the value of the determinant $\Delta = \begin{vmatrix} C & 1 & 0 \\ 1 & C & 1 \\ 6 & 1 & C \end{vmatrix}$ is:
- (1) $\frac{\sin 4\theta}{\sin \theta}$ (2) $4 \cos^2 \theta (2 \cos \theta - 1)$
 (3) $\frac{2 \sin^2 2\theta}{\sin \theta}$ (4) None of these
27. $x + ky - z = 0$, $3x - ky - z = 0$ and $x - 3y + z = 0$ has non-zero solution for k is equal to:
- (1) 0 (2) 1 (3) -1 (4) None of these
28. If $A = \begin{bmatrix} 1 & 2 \\ 3 & -5 \end{bmatrix}$, then A^{-1} is equal to:
- (1) $\begin{bmatrix} \frac{5}{11} & \frac{2}{11} \\ \frac{3}{11} & -\frac{1}{11} \end{bmatrix}$ (2) $\begin{bmatrix} -\frac{5}{11} & \frac{2}{11} \\ \frac{3}{11} & \frac{1}{11} \end{bmatrix}$ (3) $\begin{bmatrix} \frac{5}{11} & -\frac{2}{11} \\ -\frac{3}{11} & -\frac{1}{11} \end{bmatrix}$ (4) None of these
29. If $A = \begin{bmatrix} 1 & \log_b^a \\ \log_a^b & 1 \end{bmatrix}$, then $|A|$ is equal to:
- (1) 0 (2) 1 (3) \log_b^a (4) \log_a^b
30. If $y = \sqrt{\sin x + \sqrt{\sin x + \sqrt{\sin x + \dots \infty}}}$, then $\frac{dy}{dx}$ is equal to:
- (1) $-\frac{\cos x}{2y-1}$ (2) $\frac{\sin x}{1-2y}$ (3) $\frac{\cos x}{2y-1}$ (4) None of these
31. A lady gives a dinner party for six guests. The number of ways in which they may be selected from among ten friends, if two of the friends will not attend the party together, is:
- (1) 164 (2) 140 (3) 112 (4) 40
32. If w is an imaginary cube root of unity, then $(1+w-w^2)$ is equal to:
- (1) $128w$ (2) $-128w$ (3) $-128w^2$ (4) $128w^2$
33. The equation $z\bar{z} + (2-3i)z + (2+3i)\bar{z} + 4 = 0$ represents a circle of radius:
- (1) 3 (2) 4 (3) 2 (4) 6

34. If the roots of the equation $qx^2 + px + q = 0$ are complex, where p, q are real; then the roots of the equation $x^2 - 4qx + p^2 = 0$ are :
- (1) real and equal (2) imaginary
(3) real and unequal (4) none of these
35. If $a + b = 8$, then ab is greatest when :
- (1) $a = 3, b = 5$ (2) $a = 4, b = 4$ (3) $a = 6, b = 2$ (4) None of these
36. If the coefficient of 7th and 13th term in the expansion of $(1+x)^n$ are equal, then n is equal to :
- (1) 10 (2) 20 (3) 15 (4) 18
37. If ${}^n C_r$ denotes the number of combinations of n things taken r at a time, then the expression ${}^n C_{r+1} + {}^n C_{r-1} + 2 \times {}^n C_r$ equals :
- (1) ${}^{n+2} C_r$ (2) ${}^{n+2} C_{r+1}$ (3) ${}^{n+1} C_r$ (4) ${}^{n+1} C_{r+1}$
38. The number which should be added to the numbers 2, 14, 62, so that the resulting numbers may be in G. P. is :
- (1) 4 (2) 3 (3) 2 (4) 1
39. If the roots of the equation $x^3 - 12x^2 + 39x - 28 = 0$ are in A. P., then their common difference is :
- (1) ± 2 (2) ± 4 (3) ± 1 (4) ± 3
40. The equation of the straight line joining the origin to the point of intersection of $y - x + 7 = 0$ and $y + 2x - 2 = 0$ is :
- (1) $3x + 4y = 0$ (2) $4x + 3y = 0$ (3) $3x - 4y = 0$ (4) $4x - 3y = 0$
41. Let R be the relation from $A = \{2, 3, 4, 5\}$ to $B = \{3, 6, 7, 10\}$ defined by 'x divides y', then R^{-1} is equal to :
- (1) $\{(6, 2), (3, 3)\}$ (2) $\{(6, 2), (10, 2)\}$
(3) $\{(6, 2), (10, 2), (3, 3), (6, 3), (10, 5)\}$ (4) None of these
42. Which of the following is a singleton set ?
- (1) $\{x : |x| < 1, x \in \mathbb{Z}\}$ (2) $\{x : |x| = 5, x \in \mathbb{Z}\}$
(3) $\{x : x^2 = 1, x \in \mathbb{Z}\}$ (4) $\{x : x^2 + x + 1 = 0, x \in \mathbb{R}\}$

43. If $A = \{(x, y) : y = e^x, x \in R\}$ and $B = \{(x, y) : y = e^{-x}, x \in R\}$, then $A \cap B$ is :
 (1) empty set (2) not a set (3) singleton set (4) none of these
44. If $A = \{(x, y) : x^2 + y^2 = 25\}$ and $B = \{(x, y) : x^2 + 9y^2 = 144\}$, then $A \cap B$ contains :
 (1) one point (2) two points (3) three points (4) four points
45. If $z = i \log(2 - \sqrt{3})$, then $\cos z$ is equal to :
 (1) i (2) 2 (3) $3i$ (4) $2i$
46. The expression $\tan^2 \alpha + \cot^2 \alpha$ is :
 (1) ≥ 2 (2) ≤ 2 (3) ≥ -2 (4) None of these
47. A linear programming problem is concerned with finding the following value :
 (1) only maximum value (2) optimal value
 (3) only minimum value (4) none of these
48. The linear function $Z = ax + by$, where a, b are constants, which has to be maximized or minimized is called a :
 (1) constraint (2) function of any type
 (3) linear objective function (4) none of these
49. A compound statement is a statement which is made up of :
 (1) only one statement (2) any number of statements
 (3) two or more statements (4) none of these
50. A compound statement with an 'Or' is false when :
 (1) one component statement is false
 (2) none component statement is false
 (3) both the component statements are false
 (4) none of these
51. If $\vec{a} = \hat{i} + \hat{j} + \hat{k}$, $\vec{b} = 2\hat{i} - 4\hat{k}$, $\vec{c} = \hat{i} + \lambda\hat{j} + 3\hat{k}$ are coplanar, then the value of λ is :
 (1) $\frac{5}{3}$ (2) $\frac{3}{5}$ (3) $\frac{5}{2}$ (4) $\frac{7}{3}$

52. If $\vec{a} = 2\hat{i} + \hat{j} + 2\hat{k}$ and $\vec{b} = 5\hat{i} - 3\hat{j} + \hat{k}$, then the projection of \vec{b} on \vec{a} is :
 (1) 6 (2) 5 (3) 4 (4) 3
53. If $\vec{a}, \vec{b}, \vec{c}$ are mutually perpendicular unit vectors, then $|\vec{a} + \vec{b} + \vec{c}|$ is equal to :
 (1) 3 (2) $\sqrt{3}$ (3) 1 (4) 0
54. If $\vec{a} = \hat{i} + \hat{j} + \hat{k}, \vec{b} = \hat{i} + \hat{j}, \vec{c} = \hat{i}$ and $(\vec{a} \times \vec{b}) \times \vec{c} = \lambda \vec{a} + \mu \vec{b}$, then $\lambda + \mu$ is equal to :
 (1) 3 (2) 1 (3) 0 (4) 2
55. A variable plane moves, so that the sum of the reciprocals of its intercepts on the coordinates axes is $\frac{1}{2}$. Then the plane passes through :
 (1) $\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}\right)$ (2) $(-1, 1, 1)$ (3) $(2, 2, 2)$ (4) $(0, 0, 0)$
56. The equation of the plane which bisects the line joining $(2, 3, 4)$ and $(6, 7, 8)$ is :
 (1) $x + y + z - 15 = 0$ (2) $x + y + z + 15 = 0$
 (3) $x - y - z - 15 = 0$ (4) $x - y + z - 15 = 0$
57. The direction ratio of normal to the plane through $(1, 0, 0), (0, 1, 0)$ which makes an angle $\frac{\pi}{4}$ with plane $x + y = 3$ are :
 (1) $1, \sqrt{2}, 1$ (2) $1, 1, \sqrt{2}$ (3) $1, 1, 2$ (4) $\sqrt{2}, 1, 1$
58. A line makes the same angle θ , with each of the x and z axes. If the angle β which it makes with y -axis is such that $\sin^2 \beta = 3 \sin^2 \theta$, then $\cos^2 \theta$ is equal to :
 (1) $\frac{2}{5}$ (2) $\frac{1}{5}$ (3) $\frac{2}{3}$ (4) $\frac{3}{5}$
59. The solution of $\frac{dy}{dx} = 2^{y-x}$ is :
 (1) $2^x + 2^y = C$ (2) $\frac{1}{2^x} - \frac{1}{2^y} = C$ (3) $2^x - 2^y = C$ (4) None of these
60. $y + x^2 = \frac{dy}{dx}$ has the solution :
 (1) $y + x^2 + 2x + 2 = ce^x$ (2) $y + 2x = ce^x$
 (3) $y + 2x + 2 = ce^x$ (4) None of these

61. Five horses are in the race. Mr. B selects two of the horses at random and bets on them. The probability that Mr. B selected the winning horse, is :
- (1) $\frac{1}{5}$ (2) $\frac{2}{5}$ (3) $\frac{3}{5}$ (4) None of these
62. If A and B are events such that $P(A \cup B) = \frac{3}{4}$, $P(A \cap B) = \frac{1}{4}$, $P(\bar{A}) = \frac{2}{3}$, then $P(\bar{A} \cap B)$ is :
- (1) $\frac{5}{12}$ (2) $\frac{3}{8}$ (3) $\frac{5}{8}$ (4) $\frac{1}{4}$
63. A coin is tossed three times. The probability of getting head and tail alternatively, is :
- (1) $\frac{1}{8}$ (2) $\frac{1}{2}$ (3) $\frac{1}{4}$ (4) None of these
64. Seven white balls and three black balls are randomly placed in a row. The probability that no two black balls are placed adjacently, equals :
- (1) $\frac{1}{13}$ (2) $\frac{2}{15}$ (3) $\frac{7}{15}$ (4) $\frac{1}{2}$
65. The solution set of the equation $\sin^{-1} x = 2 \tan^{-1} x$ is :
- (1) $\{1, 2\}$ (2) $\{-1, 2\}$ (3) $\{-1, 1, 0\}$ (4) $\left\{1, \frac{1}{2}, 0\right\}$
66. $\tan \left[\frac{1}{2} \sin^{-1} \left(\frac{2a}{1+a^2} \right) + \frac{1}{2} \cos^{-1} \left(\frac{1-a^2}{1+a^2} \right) \right]$ is equal to :
- (1) $\frac{2a}{1+a^2}$ (2) $\frac{2a}{1-a^2}$ (3) $\frac{1-a^2}{1+a^2}$ (4) $\frac{1+a^2}{1-a^2}$
67. $\tan^{-1} \frac{x}{y} - \tan^{-1} \frac{x-y}{x+y}$; ($x > y > 0$) is equal to :
- (1) $\frac{3\pi}{4}$ (2) $-\frac{\pi}{4}$ (3) $-\frac{3\pi}{4}$ (4) $\frac{\pi}{4}$
68. If in a triangle ABC , $\angle A = \tan^{-1} 2$ and $\angle B = \tan^{-1} 3$, then angle C is equal to :
- (1) $\frac{\pi}{4}$ (2) $\frac{3\pi}{4}$ (3) $-\frac{\pi}{4}$ (4) None of these

69. For real numbers x and y , we write $xRy \Leftrightarrow x^2 - y^2 + \sqrt{3}$ is an irrational number. Then the relation R , is :
- (1) Transitive (2) Reflexive (3) Symmetric (4) None of these
70. A function f from the set of natural numbers to integers defined by
- $$f(x) = \begin{cases} \frac{n-1}{2}, & \text{when } n \text{ is odd} \\ -\frac{n}{2}, & \text{when } n \text{ is even} \end{cases} \text{ is :}$$
- (1) one-one but not onto (2) onto but not one-one
(3) one-one and onto both (4) none of these
71. If two sets A and B are having 99 elements in common, then the number of elements common to each of the sets $A \times B$ and $B \times A$ are :
- (1) 99^2 (2) 18 (3) 2^{99} (4) 100
72. If $f(x) = 1 - \frac{1}{x}$, then the value of $f\left[f\left(\frac{1}{x}\right)\right]$ is :
- (1) $\frac{1}{x} - 1$ (2) $\frac{x}{1-x}$ (3) $\frac{x-1}{x}$ (4) $\frac{x}{x-1}$
73. The function $f(x) = \log\left(x + \sqrt{x^2 + 1}\right)$ is :
- (1) an even function (2) an odd function
(3) a periodic function (4) none of these
74. The value of $\sin A \sin(60^\circ + A) \sin(60^\circ - A)$ is equal to :
- (1) $\sin 3A$ (2) $\sin \frac{3A}{2}$ (3) $\sin \frac{3A}{4}$ (4) $\sin \frac{4A}{3}$
75. If $\cos \theta = \frac{1}{2}\left(x + \frac{1}{x}\right)$, then $\frac{1}{2}\left(x^2 + \frac{1}{x^2}\right)$ is equal to :
- (1) $\cos 2\theta$ (2) $\sin 2\theta$ (3) $\sec 2\theta$ (4) $\tan 2\theta$
76. If $y = \sin^2 \theta + \operatorname{cosec}^2 \theta$, $\theta \neq 0$, then :
- (1) $y > 2$ (2) $y \leq 2$ (3) $y \geq -2$ (4) $y = 0$

77. The value of $\frac{\cos 12^\circ - \sin 12^\circ}{\cos 12^\circ + \sin 12^\circ} + \frac{\sin 147^\circ}{\cos 147^\circ}$ is equal to :
- (1) 0 (2) -1 (3) 1 (4) None of these
78. $P(n) : 1 + 3 + 5 + \dots + (2n - 1) = n^2$ is :
- (1) true for $n > 1$ (2) true for no n
(3) true for all $n \in N$ (4) none of these
79. If eleven members of a committee sit at a round table so that the President and Cashier always sit together, then the number of arrangements is :
- (1) $\underline{10} \times 2$ (2) $\underline{9} \times 2$ (3) $\underline{10}$ (4) None of these
80. In how many ways can 5 keys be put in a ring ?
- (1) $\underline{5}$ (2) $\frac{\underline{5}}{2}$ (3) $\underline{4}$ (4) $\frac{\underline{4}}{2}$
81. The straight line whose sum of the intercepts on the axes is equal to half of the product of the intercepts, passes through the point :
- (1) (2, 2) (2) (1, 1) (3) (4, 4) (4) (3, 3)
82. The equation of a circle with centre (1, 2) and tangent $x + y - 5 = 0$ is :
- (1) $x^2 + y^2 + 2x - 4y + 6 = 0$ (2) $x^2 + y^2 - 2x - 4y + 3 = 0$
(3) $x^2 + y^2 - 2x + 4y + 8 = 0$ (4) $x^2 + y^2 - 2x - 4y + 8 = 0$
83. The distance between the foci of an ellipse is 16 and the eccentricity is $\frac{1}{2}$. Length of major axis of the ellipse is :
- (1) 8 (2) 16 (3) 32 (4) 64
84. The ratio in which the line joining (2, 4, 5) and (3, 5, -4) is divided by the yz -plane is :
- (1) 4 : -3 (2) 3 : 2 (3) 2 : 3 (4) -2 : 3
85. A plane makes intercepts 3 and 4 respectively on z -axis and x -axis. If plane is parallel to y -axis, then its equation is :
- (1) $3z + 4x = 12$ (2) $3y + 4z = 12$ (3) $3x + 4z = 12$ (4) $3z + 4y = 12$

86. $\lim_{x \rightarrow 0} \frac{\int_0^x y \, dy}{x \tan(\pi + x)}$ is equal to :
- (1) 2 (2) $\frac{1}{2}$ (3) -2 (4) None of these
87. The points of discontinuity of $\tan x$ are :
- (1) $x = n\pi$ (2) $2n\pi$ (3) $(2n+1)\frac{\pi}{2}$ (4) $-2n\pi$
- where $n \in I$
88. If $xy = e^{x-y}$, then $\frac{dy}{dx}$ is equal to :
- (1) $\frac{(x-1)y}{x(1+y)}$ (2) $\frac{(x+1)y}{x(1+y)}$ (3) $\frac{(x-1)y}{x(1-y)}$ (4) None of these
89. If $y = \frac{1 + \sin x - \cos x}{1 + \sin x + \cos x}$, then $\frac{dy}{dx}$ is equal to :
- (1) $\frac{1}{\cos x}$ (2) $\frac{1}{\sin x}$ (3) $\frac{1}{1 - \cos x}$ (4) None of these
90. If $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$, then A^{100} is equal to :
- (1) $2^{100}A$ (2) $100A$ (3) $2^{99}A$ (4) $299A$
91. If $f(x) = x + 2$, then $f'[f(x)]$ at $x = 4$ is :
- (1) 8 (2) 1 (3) 4 (4) 5
92. The value of $\frac{d}{dx} \left[\left(\frac{\tan^2 2x - \tan^2 x}{1 - \tan^2 2x \tan^2 x} \right) \cot 3x \right]$ is :
- (1) $\sec x$ (2) $\sec^3 x$ (3) $\sec x \tan x$ (4) $\sec^2 x$
93. If $x = a \sin \theta$, $y = b \cos \theta$, then $\frac{d^2y}{dx^2}$ is equal to :
- (1) $\frac{b}{a^2} \sec \theta$ (2) $-\frac{b}{a^2} \sec^3 \theta$ (3) $\frac{b}{a} \sec^2 \theta$ (4) None of these

94. If $x^m y^n = (x+y)^{m+n}$, then $\frac{dy}{dx}$ is :
 (1) $\frac{y}{x}$ (2) $\frac{x}{y}$ (3) xy (4) None of these
95. Maximum slope of the curve $y = -x^3 + 3x^2 + 9x - 27$ is :
 (1) 0 (2) 16 (3) 12 (4) 32
96. The function x^x is increasing, when :
 (1) $x > \frac{1}{e}$ (2) $x < \frac{1}{e}$ (3) $x < 0$ (4) None of these
97. The rate of change of the surface area of a sphere of radius r , when the radius is increasing at the rate of 2 cm/s is proportional to :
 (1) $\frac{1}{r}$ (2) r^2 (3) r (4) $\frac{1}{r^2}$
98. Angle between the tangents to the curve $y = x^2 - 5x + 6$ at the points (2, 0) and (3, 0) is :
 (1) $\frac{\pi}{3}$ (2) $\frac{\pi}{2}$ (3) $\frac{\pi}{6}$ (4) $\frac{\pi}{4}$
99. $\int \frac{(x+1)^2}{x(x^2+1)} dx$ is equal to :
 (1) $\log x + C$ (2) $2 \tan^{-1} x + C$ (3) $\log \frac{1}{1+x^2} + C$ (4) None of these
100. $\int \frac{x + \sin x}{1 + \cos x} dx$ is equal to :
 (1) $x \tan \frac{x}{2} + C$ (2) $\tan \frac{x}{2} + C$ (3) $\log \cos \frac{x}{2}$ (4) None of these

1. 1	16. 4	31. 2	46. 1	61. 1	76. 4	91. 3
2. 4	17. 2	32. 4	47. 3	62. 4	77. 3	92. 1
3. 2	18. 3	33. 1	48. 2	63. 2	78. 4	93. 3
4. 3	19. 4	34. 3	49. 4	64. 3	79. 2	94. 4
5. 1	20. 2	35. 3	50. 1	65. 3	80. 1	95. 2
6. 1	21. 1	36. 4	51. 2	66. 1	81. 2	96. 1
7. 1	22. 2	37. 2	52. 1	67. 2	82. 1	97. 2
8. 3	23. 3	38. 1	53. 3	68. 4	83. 3	98. 3
9. 2	24. 4	39. 1	54. 3	69. 2	84. 3	99. 3
10. 4	25. 3	40. 3	55. 4	70. 1	85. 3	100. 3
11. 2	26. 2	41. 2	56. 2	71. 3	86. 2	
12. 3	27. 3	42. 4	57. 1	72. 4	87. 4	
13. 1	28. 1	43. 2	58. 2	73. 3	88. 1	
14. 3	29. 4	44. 1	59. 3	74. 1	89. 2	
15. 2	30. 3	45. 3	60. 4	75. 2	90. 3	


 02/7/16
 Professor & Head,
 Department of Mathematics
 M.D. University ROHTAK

1. 2	16. 2	31. 3	46. 1	61. 3	76. 2	91. 2
2. 4	17. 3	32. 1	47. 2	62. 4	77. 4	92. 1
3. 2	18. 1	33. 3	48. 4	63. 3	78. 1	93. 3
4. 1	19. 4	34. 4	49. 2	64. 1	79. 2	94. 3
5. 3	20. 3	35. 2	50. 1	65. 2	80. 3	95. 4
6. 1	21. 1	36. 1	51. 2	66. 4	81. 2	96. 2
7. 3	22. 4	37. 2	52. 4	67. 3	82. 3	97. 1
8. 2	23. 2	38. 3	53. 1	68. 4	83. 1	98. 2
9. 4	24. 3	39. 3	54. 3	69. 2	84. 3	99. 3
10. 1	25. 1	40. 3	55. 3	70. 1	85. 2	100. 4
11. 1	26. 1	41. 1	56. 4	71. 2	86. 4	
12. 2	27. 1	42. 4	57. 2	72. 1	87. 2	
13. 3	28. 3	43. 2	58. 1	73. 3	88. 3	
14. 4	29. 2	44. 3	59. 1	74. 3	89. 4	
15. 3	30. 4	45. 3	60. 3	75. 3	90. 2	

Handwritten signature
02/7/16
Professor & Head,
Department of Mathematics,
M.D. University, ROHTAK

1. 2	16. 1	31. 2	46. 4	61. 2	76. 1	91. 2
2. 3	17. 2	32. 1	47. 2	62. 4	77. 2	92. 1
3. 1	18. 3	33. 3	48. 1	63. 2	78. 4	93. 3
4. 3	19. 3	34. 3	49. 1	64. 1	79. 2	94. 3
5. 2	20. 3	35. 4	50. 3	65. 3	80. 1	95. 3
6. 4	21. 3	36. 2	51. 1	66. 1	81. 1	96. 2
7. 2	22. 4	37. 1	52. 2	67. 3	82. 4	97. 4
8. 3	23. 3	38. 2	53. 3	68. 2	83. 2	98. 1
9. 4	24. 1	39. 3	54. 4	69. 4	84. 3	99. 2
10. 2	25. 2	40. 4	55. 3	70. 1	85. 1	100. 3
11. 3	26. 4	41. 2	56. 2	71. 1	86. 1	
12. 1	27. 3	42. 4	57. 3	72. 4	87. 1	
13. 3	28. 4	43. 1	58. 1	73. 2	88. 3	
14. 4	29. 2	44. 3	59. 4	74. 3	89. 2	
15. 2	30. 1	45. 3	60. 3	75. 3	90. 4	

Handwritten signature
02/07/16

Department of Mathem.
M.D. University, ROHTAK

1. 3	16. 2	31. 2	46. 1	61. 2	76. 1	91. 2
2. 4	17. 1	32. 3	47. 2	62. 1	77. 1	92. 4
3. 3	18. 2	33. 1	48. 3	63. 3	78. 3	93. 2
4. 1	19. 3	34. 3	49. 3	64. 3	79. 2	94. 1
5. 2	20. 4	35. 2	50. 3	65. 3	80. 4	95. 3
6. 4	21. 2	36. 4	51. 1	66. 2	81. 1	96. 1
7. 3	22. 4	37. 2	52. 4	67. 4	82. 2	97. 3
8. 4	23. 1	38. 3	53. 2	68. 1	83. 3	98. 2
9. 2	24. 3	39. 4	54. 3	69. 2	84. 4	99. 4
10. 1	25. 3	40. 2	55. 3	70. 3	85. 3	100. 1
11. 2	26. 4	41. 3	56. 1	71. 1	86. 2	
12. 1	27. 2	42. 1	57. 2	72. 4	87. 3	
13. 3	28. 1	43. 3	58. 4	73. 2	88. 1	
14. 3	29. 1	44. 4	59. 2	74. 3	89. 4	
15. 4	30. 3	45. 2	60. 1	75. 1	90. 3	

Wander
 02/07/16
 Vinod K. Sharma,
 Department of Mathematics,
 M.D. University, ROHTA.